

DIRECTIONS FOR USE

FOLIAGE SPRAY:

For application in spring and summer any time after plants reach full leaf stage until start of dormant season. mix 3 to 5 quarts of this Brush Killer thoroughly in 100 gallons of clear water. Apply as a spray to the foliage, using sufficient pressure (80 pounds or more) and sufficient solution to assure adequate penetration and coverage of all parts of the plant. Plants which are inadequately sprayed or are tolerant should be retreated the following season.

DORMANT SPRAY:

For application to brush and trees during the dormant season (fall, and winter months), mix 1-gallon of this Brush Killer in 40 to 60 gallons of kerosene or fuel oil. Apply as a coarse spray with sufficient pressure and sufficient solution to cover fully all woody surfaces of plant, including stems, branches and twigs. If brush and trees are over 6 or 7 feet high, best results can be obtained by cutting them down and treating the stumps and sprouts.

STUMP AND BASAL SPRAY:

For application to stumps and large standing brush and trees 3 inches to 6 inches in diameter, mix 1-gallon of Brush Killer with 20 to 25 gallons of kerosene or fuel oil. Apply as a drenching spray to the top and sides of stumps. In the case of large standing brush and trees, apply as a drenching spray to the base of the plant thoroughly drenching all sides from a point 3 or 4 feet above the ground line down to the ground line. In all cases the spray must be applied until the solution begins to run off. Exposed roots must be treated. Application may be made any time of the year. Cut trees larger than 6 inches in diameter and treat stumps.

CAUTION: Harmful if swallowed. May cause skin irritation. Avoid contact with eyes, skin and clothing.

Do not use around the home, recreation areas and similar sites.

Do not graze to dairy animals within 6 weeks after application. Do not graze meat animals on treated areas within 2 weeks of slaughter.

Do not contaminate water by cleaning of equipment or disposal of wastes.

NOTE: Brush killer exposed to freezing temperature should be warmed to above 40°F. and mixed thoroughly before using.

EPA Reg. No. 7273-77

No. 438
CONCENTRATE

NET CONTENTS
GALLONS

**BRUSH
OFF**

2-2 LV BRUSH KILLER

CONTAINS

2 LBS. 2,4-D AND 2 LBS. 2,4,5-T

ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS:

*Isocetyl Ester of 2,4-Dichlorophenoxyacetic Acid.....34.5%

**Isocetyl Ester of 2,4,5-Trichlorophenoxyacetic Acid.....33.0%

INERT INGREDIENTS32.5%

*Equivalent to 22.9% of 2,4-Dichlorophenoxyacetic Acid

**Equivalent to 22.9% of 2,4,5-Trichlorophenoxyacetic Acid

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for other warnings.

MANUFACTURED BY

CROWN CHEMICALS

ROCKFORD, ILLINOIS 61101

This Brush
controls c
rows, a
higher of w
Following

Alder
Green Ash
Aspen
Barberry
River Birch
Blackberry
Huckleberry
Buckeye
Cane
Catalpa
Cedar
Cherry
Cottonwood
Custard

Tolerant a
for effecti

Do not ap
able plant
grapes, fr
soybeans,
are sensiti
spray or f
quantities

Vapors fr
plants in
should be
and from
drift. Do
water use
mounts of
porarily
growth.

Because
ing spray
such equi
ling and
Shipping
not reuse
Do not
reconditi
crushing
spill this
seeds, ins

Crown
merchant
otherwise
product
statement

No. 438
CONCENTRATE

NET CONTENTS
GALLONS



2-2 LV BRUSH KILLER

CONTAINS

2 LBS. 2,4-D AND 2 LBS. 2,4,5-T
ACID EQUIVALENT PER GALLON

ACTIVE INGREDIENTS:

*Isooctyl Ester of 2,4-Dichlorophenoxyacetic Acid.....34.5%
**Isooctyl Ester of 2,4,5-Trichlorophenoxyacetic Acid.....33.0%

INERT INGREDIENTS32.5%

*Equivalent to 22.9% of 2,4-Dichlorophenoxyacetic Acid

**Equivalent to 22.9% of 2,4,5-Trichlorophenoxyacetic Acid

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for other warnings.

MANUFACTURED BY

CROWN CHEMICALS

ROCKFORD, ILLINOIS 61101

DIRECTIONS FOR USE

This Brush Killer, properly mixed and applied, controls certain woody plants in pastures, fence rows, ditch banks, railroad and utility rights-of-way, roadside and industrial areas. Following plants are susceptible.

Alder	Dogwood	Maple
Green Ash	Elm	Mesquite
Aspen	Elm	Red Mulberry
Barberry	Gooseberry	Osage Orange
River Birch	Greenbrier	Pine
Blackberry	Wild Grape	Wild Plum
Buckbrush	Gum	Yellow Poplar
Buckeye	Hawthorn	Redbud
Cane	Hazel	Sassafras
Catalpa	Hickory	Sumac
Cedar	Honeysuckle	Sycamore
Cherry	Poison Ivy	Persimmon
Cottonwood	Junberry	Star Willow
Currant	Kudzu	Sage Brush

WARNING

Tolerant species may require repeat applications for effective control.

Do not apply to or allow contact with susceptible plants as tomatoes, beans, peas, flowers, grapes, fruit trees, ornamentals, alfalfa, clover, soybeans, tobacco or other desirable plants which are sensitive to 2,4-D or 2,4,5-T. Do not allow spray or fumes to drift onto them, since minute quantities may cause severe injury.

Vapors from this product may injure susceptible plants in the vicinity of application. Application should be carried out only when there is no hazard from drift, coarse sprays are less likely to drift. Do not contaminate irrigation water or water used for domestic purposes. Excessive amounts of this brush killer in the soil may temporarily inhibit seed germination or plant growth.

Because of the difficulty of thoroughly cleaning spraying equipment used with this product, such equipment should not be used for handling and applying other agricultural chemicals. Shipping containers should be destroyed and not reused.

Do not reuse container. Return to drum reconditioner or destroy by perforating or crushing and bury in a safe place. Do not store or spill this preparation on or near food, fertilizers, seeds, insecticides or fungicides.

NOTICE OF WARRANTY

Crown Chemicals makes no warranty of merchantability, fitness for any purpose, or otherwise, express or implied, concerning this product or its use, which extend beyond the statements on this label.