

# R M C SOLUBLE PROLIN® KILLS RATS AND MICE

## WATER SOLUBLE ANTICOAGULANT RODENTICIDE

Prolin water soluble baits are more convenient and effective because they:

1. Retard bacterial contamination.
2. Retard molding.
3. Last longer.
4. Remain acceptable to rats and mice.

This is a notable advance in anticoagulant rodenticides.

### ACTIVE INGREDIENTS:

*Sodium Warfarin, Sodium Salt of (3-(alpha-acetonylbenzyl)-4-hydroxycoumarin) .....	0.54% w/w
N <sup>1</sup> -(2-Quinoxaliny) sulfanilamide	
N <sup>1</sup> -(2-Quinoxaliny) sulfanilamide sulfaquinoxaline .....	0.50%
INERT INGREDIENTS .....	98.96%
	100.00%

### MAKES ONE GALLON

CONTENTS: 1.4108 oz. (40 grams) net weight.

\*Equivalent to 0.50% Warfarin.

†Sodium Warfarin manufactured under license from the Wisconsin Alumni Research Foundation, U. S. Patent No. 2,777,059.

### CAUTION: Keep Out of Reach of Children.

Protect pets, wildlife and domestic animals from bait. If swallowed by humans, wildlife, animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

Packed by  
**R.M.C. PRODUCTS CO.**  
Fort Dodge, Iowa 50501

Read Directions On Back Side

"EPA REGISTRATION NO. 7276-8"

### DIRECTIONS FOR USE

Dissolve contents of packet (40 grams) in 1 gallon of water with stirring. Solution will contain as active ingredients 0.0057% sodium warfarin and 0.0057% sodium sulfaquinoxaline.

**FOR RATS:** Place the solution in a 1-quart Mason jar and invert over a chicken water fount to serve as a drinking station. Set the fount on a level surface in a secluded spot where rats are accustomed to feed and where inaccessible to children, pets, wildlife, and domestic animals or in tamper proof bait boxes. Several drinking stations may be required depending upon the area of infestation.

Baiting should continue for 15 days or until complete lack of feeding is noted.

**FOR MICE:** Mice have a limited range and may not travel more than 20 feet from their nests. Therefore, it is necessary to distribute water bait in such a manner as to cover the entire infested area using several small drinking founts or shallow containers of 1/2 cup or 4 ounces liquid bait capacity, per drinking station. Set the founts on level surfaces in secluded spots where mice are accustomed to feed and where inaccessible to children, pets, wildlife, and domestic animals or in tamper proof bait boxes. Baiting should continue for 15 days or until complete lack of drinking is noted.

In the case of both rats and mice, drinking stations should not be exposed to freezing temperatures. Keep water bait containers clean and well supplied at all times.

Where a continuous source of infestation prevails from outdoors, permanent drinking stations should be used and replenished as needed.

ACCEPTED WITH COMMENTS