

# RMC

## SUPER BAR

READY FOR USE

Bars can be broken off into  
**24 — 3 1/3 oz. Bait Blocks**

Do not expose these bait blocks in areas accessible to dogs. Except when enclosed within a tamper proof bait box. "For use in wet or damp areas only" when used to control rats.

#### DIRECTIONS FOR CONTROLLING RATS:

Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

In moderately infested locations at least 2 or more blocks should be placed where rats feed, water, or travel, along walls, in corners, and concealed locations where they can feed unobserved. Each location should contain enough bait so that some will be left over the following morning.

Bait should be placed in locations not accessible to children, pets, wildlife and domestic animals, "or in tamper proof bait boxes."

Daily inspection is essential, especially in the first week, and bait supplies must be replenished as often as necessary. Under normal conditions, bait consumption will be light the first night, much heavier the 2nd and 3rd night thereafter. A gradual tapering off will occur as more and more rodents perish.

#### DIRECTIONS FOR CONTROLLING MICE

Break block in two pieces and place where mice or signs of mice have been seen. Bait stations at intervals of 8 to 10 feet are recommended. Assure an uninterrupted supply of bait for a period of not less than 15 days

**NET 2268 GRAMS (5 LBS.)**

### KILLS RATS & MICE

#### ACTIVE INGREDIENTS:

"Warfarin" "3-( $\alpha$ -acetylbenzyl)-4-hydroxycoumarin" 0.025%

"N-(2-quinoxaliny) sulfanilamide (sulfaquinoxaline)" 0.025%

INERT INGREDIENTS: 99.950%

TOTAL 100.000%

"Do not reuse container. When empty destroy by burning or discarding with trash in a safe place."

EPA REGISTRATION NUMBER 7276-11

### CAUTION:

### Keep out of reach of children.

Protect children, pets, wildlife and domestic animals from bait. If swallowed by humans, wildlife, animals or pets, this material may reduce the clotting ability of the blood and cause bleeding. In such case, intravenous and oral administration of vitamin K combined with blood transfusions are indicated as in the case of hemorrhage caused by overdoses of bis-hydroxycoumarin.

Manufactured by

**R.M.C. PRODUCTS CO.**

SATISFACTION GUARANTEED **ACCEPTED** FORT DODGE, IOWA 50501