

P 11/14

7276-11

1 of 6

1987
 632
 18
 185616

RMC Products Company, Inc.
 P.O. Box 848
 Fort Dodge, IA 50501

Gentlemen:

Subject: RMC Super Bar
 EPA Registration No. 7276-11
 Your Letter Dated November 4, 1986

You have completed all the data requirements requested under the Warfarin Standard.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), is acceptable subject to the comments listed below.

1. In the application directions for the 1-lb bar, insert hyphens between numbers and units following them. For example, change "6 1/2 oz" to "6 1/2-oz" and "8 to 12 foot" to "8 to 12-foot."
2. Under "NET WEIGHT" put English units first and metric units second.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e).

85317: Peacock: M-3: KEMCO: 2/6/87: 8/18/87: DEJ: VO: JS: TAR

SYMBOL ▶								
SURNAME ▶								
DATE ▶								

Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,



William H. Miller
Product Manager (16)
Insecticide-Rodenticide Branch
Registration Division (TS-767C)

- Enclosures: 1. Stamped Label
2. A-79 Enclosure

RMC SUPER BAR

KILLS NORWAY & ROOF RATS
AND HOUSE MICE

ACTIVE INGREDIENT:
Warfarin 3-(alpha-acetonylbenzyl)-4
hydroxycoumarin 0.025%
INERT INGREDIENTS 99.975%
TOTAL 100.000%

CAUTION KEEP OUT OF REACH OF CHILDREN
"SEE BELOW FOR ADDITIONAL CAUTION"

PRECAUTIONARY STATEMENTS HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.
NOTE TO PHYSICIAN: If ingested, administer Vitamin K¹ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD: Keep Out of Lakes, Streams or Ponds. DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamperproof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

This 16 oz. bar may be broken or cut into 6 2-1/8 oz. bars for rats. The 3 1/8 oz. bars may be broken or cut into 6 1/2 oz. or 12 1/4 oz. bars for mice.

RATS AND MICE: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water and food or feed by storage or disposal.

STORAGE: Store only in original container in a dry place.

DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

RMC PRODUCTS CO., INC.

FORT DODGE, IOWA 50501

EPA REGISTRATION NUMBER 7276-11

EPA ESTABLISHMENT NO. 7276-IA-1

NET WEIGHT 453.6 GRAMS (1 LB.)

ACCEPTED
with COMMENTS
In EPA Letter Dated:

FEB 19 1977

Under the Federal Insecticide,
Fungicide and Rodenticide Act
as amended, this pesticide
registered under EPA Reg. No.

7276-11

BEST AVAILABLE COPY

Air Back

RMC

SUPER BAR

KILLS NORWAY & ROOF RATS

Bars can be broken off into
24 — 3 1/3 oz. Bars [redacted]
NET 2268 GRAMS (5 LBS.)

ACTIVE INGREDIENT:

Warfarin 3-(alpha-acetonylbenzyl)-4 hydroxycoumarin.	0.025%
INERT INGREDIENTS	99.975%
TOTAL	100.000%

CAUTION
Keep out of reach of children.

SEE BELOW ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K¹ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times

ENVIRONMENTAL HAZARD:
Keep Out of Lakes, Streams or Ponds.

R.M.C. Products Co., Inc.
FORT DODGE, IOWA 50501

EPA Number 7276-11
EPA ESTABLISHMENT NO 7276 IA 1

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling

USE RESTRICTIONS: For control of Norway rats, and Roof rats) in [redacted] and around homes, industrial and agricultural buildings, and similar man-made structures

Treated baits must be placed in tamperproof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats [redacted] will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days

RATS [redacted] Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed

STORAGE AND DISPOSAL

Do not contaminate water and food or feed by storage or disposal.
STORAGE: Store only in original container in a dry place
DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance

KILLS NORWAY & ROOF RATS AND HOUSE MICE

ACTIVE INGREDIENT: bishydroxycoumarin 0.025%
INERT INGREDIENTS 99.975%
TOTAL 100.000%

PRECAUTIONARY STATEMENTS
HAZARD TO HUMANS AND DOMESTIC ANIMALS
Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding. If ingested, administer Vitamin K¹ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:
Keep Out of Lakes, Streams or Ponds.
DIRECTIONS FOR USE
It is a violation of federal law to use this product in a manner inconsistent with its labeling.
For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.
Treated baits must be placed in tamperproof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.
Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.


APPLICATION DIRECTIONS
Apply 4 to 16-ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.
Apply ¼ to ½-ounce of bait at 8 to 12-foot intervals. Larger placements (up to 2-ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.
This 16 oz. bar may be broken or cut into 6 3-1/5-oz. bars for rats. The 3 1/5-oz. bars may be broken or cut into 6 ½-oz. or 12 ¼-oz. bars for mice.
Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as far as possible. If reinfestation does occur, repeat treatment. Where continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL
Do not contaminate water and food or feed by storage or disposal.
STORAGE Store only in original container in a dry place.
DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinseate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

RMC PRODUCTS CO., INC.
FORT DODGE, IOWA 50501
EPA REGISTRATION NUMBER 7276-11
EPA ESTABLISHMENT NO. 7276-1A-1
NET WEIGHT 1 LB. (453.6 GRAMS)

BEST AVAILABLE COPY

NOT REVIEWED
In Accordance with FR Notice 82-2.
Based on Draft Labeling Dated



Bars can be broken off into

24 — 3 1/3 oz. Bars
NET 5 LBS. (2268 GRAMS)

ACTIVE INGREDIENT:

Warfarin (3-alpha-acetonylbenzyl)-4

hydroxycoumarin 0.025

INERT INGREDIENTS 99.975

TOTAL 100.000

CAUTION

Keep out of reach of children.

SEE BELOW ADDITIONAL PRECAUTIONARY STATEMENTS

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K¹ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

ENVIRONMENTAL HAZARD:

Keep Out of Lakes, Streams or Ponds

R.M.C. Products Co., Inc.
FORT OODGE, IOWA 50501

EPA NUMBER 7276-11

EPA ESTABLISHMENT NO. 7276-1A-1

DIRECTIONS FOR USE:

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats and Roof rats in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamperproof bait boxes or in locations *not accessible to children, pets, domestic animals or wildlife.* Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings, in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS:

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days

RATS: Replace contaminated or spoiled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present establish permanent bait stations and replenish bait as needed.

STORAGE AND DISPOSAL

Do not contaminate water and food or feed by storage or disposal.

STORAGE: Store only in original container in a dry place.

DISPOSAL: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

BEST AVAILABLE COPY