3 8 7 1 1975

RODENT BAIT

MEAT (Bits Form)

FRA PEGISTRATION NO 1216 16

12 - 5 1 3 oz. THROW PAC Enclosed

Assure an uninterrupted supply of bait for a period of not less than 10 days and continue baiting until all signs of feeding have stopped.

or 1. 3 Throw Pac of bait should be placed where rats feed water or travel along walls, in corners and concealed locations where they can feed un observed. Each location should contain enough boit so that some will be left over the following morning.

Daily inspection is essential, especially in the first week and bart supplies must be replenished as often as necessary. Under normal conditions bait consumption will be light to e first night, much enceen the 2014 and 3rd night.

First W PA meft untouched for 4 to 5 consecution in the strong be moved to a new location

The complete supply of balt for a period of not an absolute of the complete supply of balt for a period of not accomplete supply of balt for a period of not accomplete supply of balt for a period of not accomplete supply.

and the second of the second of the second

NET 1815 GRAMS (4 LBS.) KILLS NORWAY & ROOF RATS AND HOUSE MICE

Act for the type of A type & a

CAUTION.

and the second of the second

RMAN A CIAN AT ONE

Note to Physics in the second of the Physics of the

The proceed that the first of the experience of

The contraction of the contracti

Approximation of the state of t

R.M.C. PRODUCTS CO.

FORT DODGE, IOWA 50501

SATISFACTION GUARANTEED

R.M.C. RAT BAIT MAY BE USED ON FARMS, IN HOMES, MOBILE HOMES, STOCKYARDS AND PACKING PLANTS, WHEN USED AS DIRECTED.

DIRECTIONS FOR CONTROLLING NORWAY & ROOF RATS

Assure an uninterrupted supply of balt for a period of not less than 10 days and continue baiting until all signs of feeding have stopped. In moderately infested locations at least 4 to 16 ozs. or 1-3 Throw Pac of balt should be placed where rats feed, water, travel, along walls in corners and concealed locations where they can feed unobserved. Each location should contain enough balt so that some will be left over the following morning.

Daily Inspection is essential, especially in the first week and bait supplies must be replenished as often as necessary. Under normal conditions, bait consumption will be light the first night, much heavier the 2nd and 3rd nights.

Rat bait left untouched for 4 to 5 consecutive nights should be moved to a new location.

and place ¼ to ½ ozs. (tablespoon amounts) shallow containers where mice or signs of mice have been seen. Bait stations at intervals of 8 to 10 feet are recommended. Assure an un-Interrupted supply of bait for a period of not less than 15 days.

CAUTION:


RMC Meat Bait, contains as it's active ingredient an anticoagulant chemical which if taken by humans, domestic animals, of pets, may result in serious blood hemorrages. In case baits are eaten, give a teaspoonful of salt in a glass of warm water and repeat until vomit fluid is clear. CALL A PHYSICIAN AT ONCE. Note to Physician: Treat patient as for an overdose of Dicumoral. Antidote: Use of Vitamin K

Environmental Caution

Treated balts should be placed in locations not accessible to children, pets, wildlife and domestic animals, or in tamperproof balt boxes.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.



RMC PRODUCTS CO., FORT DODGE, IO NA EPA ESTABLISHMENT NO. 7276-IA-1