

PM14

7276-17

1 of 1

REREGISTERED

ACCEPTED

JUN 9 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under 7276-17

RMC

MEAT BAIT

SOFT PELLETS

KILLS NORWAY & ROOF RATS and HOUSE MICE

4 1/2 Bags Enclosed

NET 1815 GRAMS (4 LBS.)

ACTIVE INGREDIENTS

Warfarin (3-(4-chlorophenyl)-4-hydroxycoumarin)	0.025%
INERT INGREDIENTS	99.975%
TOTAL	100.000%

CAUTION

Keep out of reach of children.

SEE RIGHT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

R.M.C. Products Co., Inc.
FORT DODGE, IOWA 50501

EPA REGISTRATION NO. 7276-17
EPA ESTABLISHMENT NO. 7276-1A-1

ENVIRONMENTAL HAZARD:
Keep Out of Lakes, Streams or Ponds.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Keep away from humans, domestic animals and pets. If swallowed, this material may reduce the clotting ability of the blood and cause bleeding.

NOTE TO PHYSICIAN: If ingested, administer Vitamin K₁ intramuscularly or orally, as indicated in bishydroxycoumarin overdose. Repeat as necessary based on monitoring of prothrombin times.

HAZARD TO HUMANS (CHILDREN)

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

USE RESTRICTIONS: For control of Norway rats, Roof rats and House mice in and around homes, industrial and agricultural buildings, and similar man-made structures.

Treated baits must be placed in tamperproof bait boxes or in locations not accessible to children, pets, domestic animals or wildlife. Do not place bait in areas where there is a possibility of contaminating food or surfaces that come in direct contact with food.

SELECTION OF TREATMENT AREAS: Determine areas where rats and/or mice will most likely find and consume the bait. Generally, these areas are along walls, by gnawed openings in or beside burrows, in corners and concealed places, between floors and walls, or in locations where rodents or their signs have been observed. Protect bait from rain or snow. Remove as much food as possible.

APPLICATION DIRECTIONS

RATS: Apply 4 to 16 ounces of bait per placement. Maintain an uninterrupted supply of fresh bait for at least 10 days.

MICE: Apply 1/4 to 1/2 ounce (1-2 level tablespoons) of bait at 8 to 12 foot intervals. Larger placements (up to 2 ounces) may be needed at points of very high mouse activity. Maintain an uninterrupted supply of fresh bait for at least 15 days.

RATS & MICE: Replace contaminated or trampled bait immediately. Collect and dispose of all dead animals and unconsumed bait properly. To prevent reinfestation, eliminate food, water, and harborage as much as possible. If reinfestation does occur, repeat treatment. Where a continuous source of infestation is present, establish permanent bait stations and replenish bait as needed.

DISPOSAL

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.