


7583-4

7583-4



CUSTOMER'S PROOF

DIRECTIONS FOR USE FLYING INSECTS

Exposed stages of Flies, Mosquitoes, Gnats, Bees, Wasps, Hornets, Indian Meal Moths, Angoumois Grain Moths, Mediterranean Flour Moths, Fruit Flies, Cigarette Beetles, Flying Clothes Moths. To control these and similar insects fog this material at the rate of 1 ounce per 1000 cubic feet. Turn off all ventilating equipment, close all windows and doors. Keep treated area closed for at least 10 minutes. For small flying insects (flies, mosquitoes, gnats) close up the infested area and fill it with mist. It is frequently desirable to sweep up and destroy the fallen insects at the end of the 10 minute period.

CRAWLING INSECTS

Exposed stages of Cockroaches, Waterbugs, Silverfish, Spider Beetles, Meal Worms, Caddises, Confused Flour Beetles, Saw-Toothed Grain Beetles, Spiders, Weevils, Dermestids.

For space spray-fog 55 at 2 ounces per 1000 cubic feet. As a contact spray apply 55 directly on insects or wherever they are seen. Repeat as necessary. For Cockroaches, Waterbugs and Silverfish—thoroughly and repeatedly spray infested area, directing spray into cracks and other hiding places. Try to hit as many insects as possible

Net Contents U.S. Gallons

FIFTY-FIVE

ACTIVE INGREDIENTS:	BY WEIGHT
Petroleum Hydrocarbons	98.60%
*Technical Piperonyl Butoxide	0.50%
**N-Octyl Bicycloheptene Dicarboximide	0.50%
Pyrethrins	0.40%
*Equivalent to 0.40% of (Butylcarbityl) (6-Propylpiperonyl) Ether and 0.10% Related Compounds.	
**U.S. Patent No. 2,476,512	

CAUTION

**Keep Out of Reach of Children
Additional Cautions on Side Panel**

U.S.D.A. REGISTRATION NO. 7583 -- 4

FOR USE IN: Food Processing Plants, Restaurants, Bottling Plants, Institutions, Warehouses, Offices, Motels, Packing Houses, Canneries, Hospitals, Bakeries, Markets, Locker Rooms, Hotels, Stores, Theatres, Laundries, Box Cars, Kennels, Breweries.

AERO CHEMICAL CORPORATION

CHICAGO, ILLINOIS

during spraying. Repeated applications are necessary for control. For Flour and Grain Pests clean all farinaceous material out of bins, elevator boots and spouts and apply 55. Apply 55 as a coarse wet spray at the rate of one gallon per 1000 square feet or to the point of run-off using a pressurized sprayer. Thoroughly treat all harborages and breeding areas, cracks, crevices, under sides of work tables, behind sinks, walls, window and door frames, elevator shafts, and pits behind and under equipment, laundry carts, garbage areas, warm dark moist areas, void spaces and shelves. Repeat as necessary.

POISON

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Do not take internally. Avoid contamination of food. Food processing plants should be fogged when not in operation. Food should be removed or covered during treatment. Food processing equipment should be covered or thoroughly cleaned before re use after treatment. Do not remain in treated areas and ventilate the area after treatment is completed.

STORE AWAY FROM HEAT OR OPEN FLAME

Soiled or dusty fabrics may be stained by the fog.