## **USE DIRECTIONS:**

#### Disinfection

Use at 2 ounces per gallon. This product will be effective against Pseudomonas aeruginosa PRD-10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use Tar-Sar Plus with a cloth or mop and apply to walls, floors and other hard surfaces.

For heavy soil or organic matter, a pre-cleaning step is recommended.

#### Sanitization

If used for sanitization of previously cleaned food equipment or food contact items, limit the level of active quaternaries to 200 ppm (2.8 ounces per 5 gallons of water). NO RINSE REQUIRED.

This product when used on environmental, inanimate, hard surfaces at 2 ounces per gallon of water, is effective against Influenza A<sub>2</sub>, Herpes Simplex, Adenovirus type 2, and Vaccinia viruses.

### PHENOL COEFFICIENTS:

HENGE COLLINOS.	
Guaranteed	Minimu
S. aureus	44
S. typhosa	28

Rinse empty container thoroughly with water and discard.


### CLEANER DEODORIZER

# TAC-SAN PLUS DISINFECTANT FUNGICIDE

Designed for Hospital and Institutional use.

NET CONTENTS: \_\_\_\_ EPA Reg No. 7350-18

# WARNING-KEEP OUT OF REACH OF CHILDREN. SEE RIGHT SIDE PANEL FOR ADDITIONAL PRECAUTIONARY AND FIRST AID STATEMENTS.



# CHASKA CHENICAL

304 MASTER AVENUE SAVAGE, MINN. 55378 PHONE 612/890-1820

APROMETED

APROMETED

WHOLE AND RODENTICIDE ACT

FOR EGONGMIC POISON REGISTER
ED UNDER NOUTS OF SUBJECT

TO ATTACHED COMMENTS

Tac-San Plus is a cleaner, disinfectant, and santizer, designed for institutional use such or Hospitals, Nursing Homes and Institutions where house keeping will be of prime importance in controlling thhazard of cross-infection from environmental surfaces. Tac-San Plus can provide the assurance you need in this area, when used as directed. For use on floors walls, sink tops, garbage pails, telephones, restrooms, bed pans and operating rooms. Effective against Staphylococcus aureus, Salmonella choleraesuis and Pseudomonas aeruginosa. Also effective against pathogenic fungi, Trichophyton interdigitale.

When used for sanitization of previously cleaned food processing equipment and food contact items limit active quaternaries to 200 ppm. At this level NO POTABLE WATER RINSE is required.

### WARNING

Keep Out of Reach of Children. Causes eye and skin irritation. Do not get in eyes, on skin or or clothing. Harmful if swallowed. Avoid contamination of food.

### FIRST AID

In case of contact, immediately flush eyes or skir with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash all contaminater clothing before reuse. If swallowed, drink milk, eg whites, gelatin solution, or if these are not available drink large quantities of water. Call a physician.