

GEORGIA-PACIFIC SPLASHER POOL LIQUID SANITIZER is a specially designed product for controlling algae and bacteria in pools up to about 5,000 gallons. Germs can be transmitted even in freshly filled pools

when there is no sanitizer present, so it is important to treat even pools which are emptied after each use. Pools which are not emptied after each use must be treated to prevent algae and disinfect against bacteria.

DIRECTIONS FOR USE (Swimming Pools)

- A. Pools which are emptied after each use.
1. Calculate approximate pool capacity, using chart below.
 2. Add one (1) fluid ounce Splasher Pool Liquid Sanitizer for each 100 gallons of pool capacity. Blend this in well with the pool water, preferably by adding during filling of the pool. Check residual chlorine with a reliable test kit. If below 1.0 part per million (PPM), repeat dosage until a chlorine residual of 1.0 PPM is established.
 3. As long as pool is in use, test pool water frequently with the test kit and add product as needed to maintain 1.0-1.5 PPM residual chlorine.
- B. Pools which are not emptied after each use.
1. Follow first two steps above, except add product by broadcasting over pool surface in several locations.
 2. Measure pH immediately and at least once each week with a reliable pool test kit. Adjust pH to 7.2-7.8 if necessary by adding G-P Liquid Acid or G-P Dry Acid to reduce pH, or by adding G-P Soda Ash to increase pH.*
 3. Maintain residual chlorine by adding one (1) fluid ounce Splasher Pool Liquid Sanitizer for each 100 gallons pool capacity every other day or as needed to maintain 1.0-1.5 PPM residual chlorine.

*Follow Directions for Use on Labels of these Products.

HOW TO CALCULATE POOL CAPACITY

Shape of Pool	Approximate Gallons of Water (Dimensions in Feet)
Rectangular	Average depth x average length x average width x 8
Circular	Diameter x diameter x average depth x 6
Oval with straight sides	Full width x full length x average depth x 7

IMPORTANT NOTES:

It is good practice to "superchlorinate" pools not emptied after each use each 2-4 weeks. To superchlorinate, add two (2) ounces Splasher Pool Liquid Sanitizer per 100 gallons pool capacity. Resume regular program when residual chlorine falls to 1.0-1.5 PPM.

For pools which are not emptied after each use, of any size, it is extremely important to use a test kit for available chlorine and pH determination. These factors cannot be measured by merely looking at the pools, and if not within recommended limits, your pool may not only develop algae but can transmit germs as well.

CAUTION: Harmful if swallowed. Keep out of reach of children. May cause severe eye and skin irritation. Add only to water; do not mix with other swimming pool chemicals or any other products. Fire, explosion, or evolution of toxic gases may result.

Rinse empty container thoroughly with water and discard it.

FIRST AID: If swallowed, feed milk, egg whites, or large quantities of water. If splashed in eyes, flood immediately with water. If in contact with skin, flush with plenty of water. Call a doctor right away.

ENVIRONMENTAL CAUTIONS: This product is toxic to fish. Treated effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Another Quality Product From

Georgia-Pacific  Los Angeles, California 90040

EPA Reg. No. 7368-31-AA

EPA Est. No. 7368

ACCEPTED
368-31
AUG 1 1975
BY THE FEDERAL INSECTICIDE
AND RODENTICIDE ACT
ECONOMIC POISON REGISTER
NUMBER NO.

COMMENT

ACCEPTED WITH COMMENTS