

# LEMON SANI-CLEAN DETERGENT SANITIZER

PM 31 7368-56 11 27 90

DISINFECTANT-CLEANER-SANITIZER  
FUNGICIDE-DEODORANT

ACTIVE INGREDIENTS  
n-ALKYL ISO-C14-40% C12-10% C16-DIMETHYL  
BENZYL AMMONIUM CHLORIDE 4.8%  
INERT INGREDIENTS 95.2%

In accordance with

Phase 2 Phenol Coeff. S aureus 35  
Styphosa 47

100% DISINFECTANT

**KEEP OUT OF REACH OF CHILDREN**

## DANGER

Corrosive. See antidote statement and additional precautions on back panel. Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

### NET CONTENTS 1 GALLON (3.785 LITERS)

## Grow Group, Inc.

2501 Main Avenue

City of Commerce, CA 90040

SANI-CLEAN is an effective anti-bacterial cleaner for use in household, industrial or institutional cleaning. When used as directed, SANI-CLEAN will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial, anti-fungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

#### FOR CLEANING

To clean toilet areas, surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of SANI-CLEAN per gallon of water. Floor cleaned area with hot water. Heavily soiled floors, mops or brush used surface with a solution containing 2 ounces of SANI-CLEAN per gallon of water. Rinse cleaned surface with fresh water.

#### GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, rest rooms and other public areas, the porous surface to be treated with a solution containing 2 ounces of SANI-CLEAN per gallon of water. Surfaces that are heavily soiled should be cleaned with a brush. If SANI-CLEAN per gallon of water and then with clean water. Surfaces which may later come in contact with food must be rinsed with clean water of drinking quality. To disinfect walls, shower stalls, washrooms, rest rooms and other areas, use a spray bottle containing a solution containing 2 ounces per gallon of water. The solution will destroy the bacterium, tubercle, and a causative agent of streptococci.

#### SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first, clean thoroughly with a solution containing 1 ounce of SANI-CLEAN per gallon of water. If surface is heavily soiled, use SANI-CLEAN at a rate of 2-3 ounces per gallon to insure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1/2 ounce of SANI-CLEAN per gallon of water. All surfaces should be exposed to the sanitizing solution for a period of not less than 1 minute. Drain surfaces and allow to air dry. Do not rinse.

#### HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of SANI-CLEAN per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

#### DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of SANI-CLEAN per gallon of water.

#### DANGER

**KEEP OUT OF REACH OF CHILDREN.** This product causes eye damage and skin irritation. Do not get in eyes, on skin or on anything to be eaten, and skin when handling. Harmful or fatal if swallowed. Avoid contact with food.

#### FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of fresh water. Get medical attention if these are not available, drink large quantities of water until relieved. Call a physician immediately.

#### NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

EPA Reg. No. 7368-B6

FPA Reg. No. 7368-1x1