

LEMON SANI-CLEAN DETERGENT SANITIZER

DISINFECTANT-CLEANER-SANITIZER
FUNGICIDE-DEODORANT

BEST AVAILABLE COPY

ACTIVE INGREDIENTS
n-ALYL (50%AC) 40%AC 12, 10%AC 18 DIMETHYL
BENZYL AMMONIUM CHLORIDE

48%

INERT INGREDIENTS
Phenol Coat S surfactant—35
S hypodermic—42

95.2%

Not Reviewed. Registrant claims to be in accordance with

KEEP OUT OF REACH OF CHILDREN

DANGER

See antidote statement and additional precautions on side panel. Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

NET CONTENTS 1 GALLON (3.785 LITERS)

Grow Group, Inc.

2501 Malt Avenue

City of Commerce, CA 90040

SANI-CLEAN is an effective anti-bacterial cleaner designed for use in household, industrial or institutional cleaning. When used as directed, SANI-CLEAN will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial antifungal detergent in homes, schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired. See left panel for use directions.

FOR CLEANING

Lightly soiled areas—surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of SANI-CLEAN per gallon of water. Rinse cleaned area with fresh water. Heavily soiled areas—mop or brush soiled surface with a solution containing 2–3 ounces of SANI-CLEAN per gallon of water. Rinse cleaned surface with fresh water.

GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in homes, schools, restaurants, restrooms and other public areas, mop or brush surface to be treated with a solution containing 1 1/2 ounces of SANI-CLEAN per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of SANI-CLEAN per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality.

To disinfect locker rooms, shower stalls, walkways around pools and other areas against pathogenic fungi, mop area with a solution containing 2 ounces in 3 gallons of water. This solution will destroy Trichophyton interdigitale, a causative agent of athlete's foot.

SANITIZATION

To sanitize walls, floors and equipment in food processing plants, first clean thoroughly with a solution containing 1 ounce of SANI-CLEAN per gallon of water. If surface is heavily soiled, use SANI-CLEAN at a rate of 2–3 ounces per gallon to ensure thorough cleaning. Sanitize cleaned surface with a rinse solution containing 1/2 ounce of SANI-CLEAN per gallon of water. All surfaces should be exposed to the sanitizing solution for a period not less than 1 minute. Drain surfaces and allow to air dry. Do not rinse.

HOSPITAL DISINFECTION

To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 1/2 ounces of SANI-CLEAN per gallon of water. Those surfaces that are heavily soiled with blood, fecal material or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water and then rinsed with a fresh solution containing 2 ounces per gallon of water before disinfection.

DEODORIZATION

To deodorize surfaces during cleaning, mop or brush surfaces with a solution containing 2 ounces of SANI-CLEAN per gallon of water.

DANGER

KEEP OUT OF REACH OF CHILDREN. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful or fatal if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gastric solution, etc. If these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

EPA Reg. No. 7368-86
EPA Est. No. 7368-TX-1

Grow Group, Inc.

2501 Malt Avenue

City of Commerce, CA 90040

BEST AVAILABLE COPY