

USE DIRECTIONS

Use at 2 ounces per gallon. This product will be effective against *Pseudomonas aeruginosa* PRD 10 at this level. This has been substantiated by the AOAC Use Dilution Confirmation Test. At this level, the product is fungicidal against pathogenic fungi.

For best results, use H-D-S with a cloth or mop and apply to walls, floors and other hard surfaces.

If used for sanitization of food equipment or food contact items, limit the level of active quaternary to 200 ppm (2.8 ounces per 5 gallons).

NO RINSE REQUIRED.

For heavy soil or organic matter, a precleaning step is recommended.

Rinse empty container thoroughly with water and discard.

E.P.A. Registration Number 7389-6

Net contents on Container

Hospital DEO-SAN

Detergent-Deodorant-Sanitizer-Disinfectant-Fungicide

ACTIVE INGREDIENTS:

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	2.25%
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	2.25%
Sodium Carbonate	3.00%
Essential Oils	.50%
INERT INGREDIENTS	92.00%
	100.00%

Minimum Phenol Coefficients AOAC

C. Staphylococcus Aureus	42
Salmonella Typhosa	27

DANGER Keep Out of Reach of Children.

See Antidote/First Aid Statement on right panel. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful or fatal if swallowed. Avoid contamination of food.

NOT A HOUSEHOLD PRODUCT
FOR PROFESSIONAL HOSPITAL INSTITUTIONAL USE

Manufactured by

Maintenance Products, Inc.

374 S. FRONT STREET
MEMPHIS, TENNESSEE 38103

H-D-S is a cleaner, disinfectant, and sanitizer, designed for institutional use such as Hospitals, Nursing Homes and institutions where housekeeping will be of prime importance in reducing the hazard of cross-infection. H-D-S can provide the assurance you need in this area, when used as directed. For use on floors, walls, sink tops, garbage pails, telephones, rest rooms, bed pans and operating rooms. Effective against *Staphylococcus aureus*, *Salmonella choleraesuis* and *Pseudomonas aeruginosa*. Also effective against pathogenic fungi, *Trichophyton interdigitale*.

When used for sanitization of food processing equipment and food contact items, limit active quaternary to 200 ppm. At this level, **No Potable Water Rinse** is required.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.