

6-23-67

7393-9

ED UNDER NO.

CONTENTS 1 PINT

~~base worm, cabbage looper, caterpillars, flea beetles, leafhoppers, mites~~, apply 2 teaspoonfuls of 50% Malathion in 1 gal. of water, or 2 tablespoonfuls in 3 gal. or 1 pint in 50 gallons of water. Make full coverage application and repeat as necessary. Do not apply within 7 days of harvest.

~~Okra, onions and peppers~~ to control aphids, Japanese beetle, thrips, and pepper maggots apply 1 tablespoonful of 50% Malathion in 1 gallon of water or 1 pint in 32 gallons of water. Cover plants thoroughly. Do not apply on okra after pods have started to form. On onions and peppers do not apply within 3 days of harvest.

~~Tomatoes~~: For control of spider mites, aphids, and tomato russet mite apply 1 1/2 teaspoonful in 1 gallon of water or 1 tablespoonful in 2 gallons of water or 1 pint in 64 gallons of water. For Drosophila use 1 tablespoonful per gallon of water, or 1 pint in 32 gallons of water. Make full coverage application on foliage and fruit. Allow 1 day between spraying and harvest.

~~Fleas: Dogs and Cats~~: Spray area with a mixture of 1/2 pint 50% Malathion in 8 gallons of water. Be sure to cover thoroughly and at the same time treat dogs with same mixture, or 2 tablespoons to 1 gallon of water and spray dog thoroughly.

~~Houseflies: To help control the nuisance of houseflies around buildings, for domestic animals and around outside of homes: Mix 1/4 pint (8 tablespoons) to 1 gallon of water. Adding 1/2 cup (8 tablespoons) of sugar to mixture will help.~~

**CAUTION:** Harmful if inhaled or swallowed. Avoid breathing spray mist. Avoid repeated or prolonged contact with skin. Wash thoroughly with soap and water after using. Avoid contamination of feed and foodstuffs. Keep out of reach of children. Keep container closed and away from heat, sparks, and open flame.

WONDER CHEMICAL CO. INC.,  
SAN ANTONIO, TEXAS

USDA REG. NO. 7393-9

## FOUR SEASONS

# 50% MALATHION

Use for the control of insects on pecans, fruit trees, vegetables, ornamentals and flowers and for the control of houseflies and fleas.

#### ACTIVE INGREDIENTS:

Malathion\* . . . . . 50%

Xylene Range Aromatic

Hydrocarbon . . . . . 42.3%

INERT INGREDIENTS . . . . . 7.7%

100.0%

\*O, O-dimethyl dithiophosphate of diethyl mercaptosuccinate.

**WARNING: KEEP OUT OF REACH OF CHILDREN**  
SEE SIDE PANELS FOR ADDITIONAL CAUTIONS

#### DIRECTIONS:

Read entire label and use in accordance with label directions. Do not use on Boston, Maiden-hair and Pteris Ferns, and some species of Crassula since plant injury may result.

**PECANS:** Case bearers, aphids, pecan phylloxera, spider mites and spittle bugs. Use 2 teaspoons of 4 SEASONS 50% Malathion per gallon of water, 2 tablespoons per 3 gallons of water, or 1/2 pint per 25 gallons of water. Spray thoroughly and repeat in 7 to 10 days if necessary.

#### ORNAMENTAL SHRUBS, FLOWERS AND SHADE TREES:

For scale insects such as azalea scale, black scale crawlers, euonymus scale, Fletcher scale, juniper scale, leaf scale, magnolia scale, oyster shell scale, scurfy scale and soft brown scale use 1 tablespoonful per gallon of water. For aphids use 2 teaspoons per gallon of water. To control red spider mites, spruce mites, woolly aphid, birch leaf miners, flour-lined leaf bug, Japanese beetle adults, lace bugs, leafhoppers, mealybugs, tarnished plant bugs, thrips, whitefly and European pine shoot moth use 1 tablespoonful per gallon of water.

**Fruit Trees:** Apples, grapes, pears, apricots, plum, prune, peaches. To control aphids use 2 teaspoons per gallon of water. For red spider mites, mealybugs, white flies, woolly aphids, thrips, Japanese beetle adults, tarnished plant bugs, use 1 tablespoonful per gal. water. For scale crawlers refer to ornamentals. Malathion may cause injury to Bosc pears, Ribier grapes, McIntosh and Courtland apple varieties. Do not apply within 7 days of harvest.

**VEGETABLES:** Beans, cucumber, squash, potatoes and melons to control aphids, spider mites, leafhoppers, Mexican bean beetles, cucumber beetles, pickleworms and squash vine borer use 1 1/2 teaspoonful in 1 gallon of water or 1 tablespoonful in 2 gallons of water or 1 pint in 64 gallons of water. Be sure to get good coverage. Repeat as necessary. Do not apply to curcurbits unless plants are dry and allow 1 (one) day between application and harvest. Cabbage, turnips, broccoli, mustard greens, carrots, cauliflower, head lettuce, radishes, parsnips, spinach: to control aphids imported car-

7393-9



Okra, onions and peppers  
are planted in the same row.

Okra, onions and peppers  
are planted in the same row.

Tomatoes  
are planted in the same row.

Fig, Date and Eggplant  
are planted in the same row.

Housefly  
are planted in the same row.

CAUTION  
Do not plant in the same row.

PEACHES  
are planted in the same row.

ORNAMENTAL SHRUBS, FLOWERS AND SHADE  
TREES

Planting in the same row.

Fruit Trees  
are planted in the same row.

ORNAMENTAL SHRUBS, FLOWERS AND SHADE  
TREES