

Okra: corn earworm and stink bugs.

Lettuce and Carrots: flea beetles, harlequin bug and leafhoppers, six-spotted leafhopper (aster yellows vector), armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs and stink bugs.

Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi: flea beetles, harlequin bug, armyworms, cabbage caterpillars and corn earworm.

Ornamental Insect Control: All from Abelia and Ash to Zinnia: for control of aphids, bagworm, leaf miner, blister beetles, boxelder bug, boxwood leaf miner, flea beetles, gypsy moth, Japanese beetle, June beetle, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, rose slug, scale insects, tent caterpillars, thorn bugs, thrips (exposed).

LAWN INSECTS: Use 1 level cupful in 15 to 20 gallons of water for each 500 square feet for known control of: ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworms (lawn moths) and millipedes. For best results mow lawn and apply after watering or rain. Repeat in 2-3 weeks if necessary.

Does not control spider mites, but is compatible with all common miticides.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid prolonged or repeated contact with skin. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptacles. Do not store near feed and food products. Wash thoroughly after using.

CAUTION: For protection of honey bees avoid use if possible during periods when honey bees are visiting the crop. Warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

USDA Reg. No. 7393-15

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 50%

INERT INGREDIENTS: 50%

U. S. PATENT NO. 2,903,478.
*TRADEMARK OF UNION CARBIDE CORP.

ONE POUND

FOUR SEASONS

50W SEVIN INSECTICIDE SPRAY

For The Control of Insect Pests

PRE-HARVEST USE LIMITATION

NO LIMITATION on Beans, Blueberries, Carrots, Cucumbers, Eggplants, Grapes, Melons, Peppers, Pumpkins, Squash, Tomatoes, and Okra.

ALLOW 1 DAY between spraying and harvest of: Apples, Cherries, Cranberries, Peaches, Pears, Plums and Strawberries.

ALLOW 3 DAYS between spraying and harvest of: Cabbage, Cauliflower, Kohlrabi, Lettuce, Brussels Sprouts, Broccoli, Nectarines and Apricots.

ALLOW 5 DAYS between spraying and harvest of: Citrus fruit.

DO NOT USE on Pecans after husks split.

WARNING. KEEP OUT OF REACH OF CHILDREN
SEE OTHER CAUTIONS ON SIDE PANEL

WONDER CHEMICAL CO., INC., San Antonio, Texas

7393-15

DIRECTIONS: Read Entire Label.
Use strictly in accordance with directions and cautions.

REGULAR MIXTURE 2 Level Tablespoons per Gallon of Water.

Be sure you are using sufficient spray gallonage to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary.

Some plants and the insects controlled on them are:

Apples and Pears: Apple sucker, green apple aphid, wooly apple aphid, bagworm, California pear-slug, codling moth, apple rust mite, pear leaf blister mite, pear rust mite, pear psylla, teniform leaf miners, San Jose scale. Do not use until 30 days after full bloom.

Peaches, Apricots and Nectarines: codling moth, cucumber beetles, European earwig, Japanese beetle, Pandemis moth, June beetles, orange tortrix, oriental fruit moth, peach twig borer, periodical cicada, plum curculio, fruit tree leaf roller, red-banded leaf roller, Platynota flavendana, tussock moth, Lecanium scales, San Jose scale and olive scale.

Plums, Prunes, and Cherries: black cherry aphid, mealy plum aphid, cherry fruit fly, cherry fruitworm, eye-spotted bud moth, fruit tree leaf roller, red-banded leaf roller, Japanese beetle, peach twig borer, plum curculio, prune leafhopper, brown soft scale, Forbes scale, Lecanium scales, oystershell scale and San Jose scale.

Grapes: European fruit lecanium, grape leaf folder, grape leafhopper, and grape leaf skeletonizer.

Strawberries: meadow spittlebug, strawberry leaf roller and strawberry weevil.

Blueberries: blueberry maggot, cherry and cranberry fruitworms, European fruit lecanium and Japanese beetle.

Cranberries: cutworms, fireworms, fruitworms, Japanese beetle and leafhoppers.

Pecan: pecan nut casebearer.

Beans: (Green beans, lima, snap cowpeas and black-eyed peas), bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar and western bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bug.

Cucumber, Melons, Pumpkin and Squash: Pickle worm, melonworm, cucumber beetles, flea beetles leafhoppers and squash bugs.

Potato, Tomato, Eggplant and Pepper: Colorado potato beetle, flea beetle and leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

FOUR SEASONS

WINTER

SPRING

SUMMER

FALL

WINTER

SPRING

SUMMER

FALL

WINTER

SPRING

WINTER

SPRING

SUMMER

FALL

WINTER

SPRING

SUMMER

FALL

WINTER

SPRING

SUMMER

FALL