

ACCEPTED

Feb 16 1971

Cabbage, Broccoli, Brussels Sprouts, and Cauliflower: Flea beetles, harlequin bug, armyworms, cabbage caterpillars and corn earworm.

Dogs and Cats: Ticks and Fleas. Dust liberally over entire animal, avoiding eyes; rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Also, dust liberally in and around sleeping quarters, kennels and dog houses. Be sure to force dust into cracks and crevices. Repeat as necessary. Avoid treatment of kittens and puppies less than 4 weeks old.

Lawn Insects: Dust liberally using 1 lb. per 275 sq. ft. for control of ants, chinch bugs, earwigs, fall armyworm, fleas, leafhoppers, mosquitoes, sod webworms (lawn moths), millipedes. For best results, mow lawn and apply after watering or rain. Repeat in 2-3 weeks if necessary.

Does not control spider mites but is compatible with all common miticides.

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs, and watering receptacles.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops. Protective information may be obtained from your Co-operative Agricultural Extension Service.

ACTIVE INGREDIENTS

- Carbaryl (1-naphthyl)
- N-methylcarbamate

INERT INGREDIENTS

Trademark of Cyanamid Corporation
© 1971, Cyanamid Corporation, New York, N.Y.
EPA Reg. No. 101-147
USDA REG. NO. 749116



FOR THE CONTROL OF INSECT PESTS

Pre-harvest use Limitation

NO LIMITATION on Beans, Blueberries, Carrots, Cucumbers, Eggplants, Melons, Grapes, Peppers, Pumpkins, Squash, Tomatoes, and Okra.

ALLOW 1 DAY between dusting and harvest of: Cranberries and Strawberries.

ALLOW 3 DAYS between spraying and harvest of: Cabbage, Cauliflower, Brussels Sprouts, and Broccoli.



KEEP OUT OF REACH OF CHILDREN

WONDER CHEMICAL CO., INC. SAN ANTONIO, TEXAS 78201

Application Rate

Use sufficient dust to obtain full coverage. Apply when insects or their damage appear. Repeat at 7 to 10 day intervals or as necessary. Some plants and the insects controlled on them are:

Grapes: grape leaf folder, grape leafhopper and grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles and red banded leaf roller.

Strawberries: for control of meadow spittlebug, strawberry leaf roller and strawberry weevil.

Blueberries: blueberry maggot, cherry and cranberry fruitworms and Japanese beetles.

Cranberries: cut worms, fireworms, fruitworm, Japanese beetle and leafhoppers.

Beans: (Green, Lima, Snap beans, cowpeas and blackeyed peas) Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar, and western bean cutworm, armyworms, corn earworm, stink bugs and tarnish plant bug.

Cucumber, Melons, Pumpkin and Squash: pickleworm, rootworm, cucumber beetles, flea beetles, leafhoppers, and squash bugs. May cause some leaf injury due to prolonged high humidity. Do not use on watermelons, Florida.

Tomato, Eggplant and Pepper: Colorado potato beetle, flea beetles and leafhoppers. European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tomato pinworm, and stink bugs.

Okra: corn earworm and fall armyworm. Repeat 7 to 10 day intervals.

Carrots: flea beetle, root fly, root knot, root hoppers, six spotted leafhopper, carrot root fly, carrot rootworm, cabbage caterpillar, carrot root fly, meadow spittlebug, and carrot root fly.

FOR CONTROL OF CERTAIN VEGETABLE AND CERTAIN FRUIT INSECT PESTS

Does not control spider mites but is compatible with all common miticides.

CAUTION.

KEEP OUT OF REACH OF CHILDREN

SEE OTHER CAUTIONS ON LAW

Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs, and watering receptacles.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

ACTIVE INGREDIENTS

Carbaryl (1-naphthyl N-methyl carbamate) 5%

INERT INGREDIENTS 95%

NET WT. 5 LBS.



Manufactured by Wonder Chemical Co., Inc.
San Antonio, Texas 78201
EPA Reg. No. 105-102-01

WONDER CHEMICAL CO., INC., SAN ANTONIO, TEXAS 78201

Grapes: grape leaf folder, grape leafhopper and grape leaf skeletonizer, grape berry moth, Japanese beetle, June beetles and red-banded leaf roller.

Strawberries: for control of meadow spittlebug, strawberry leaf roller and strawberry weevil.

Blueberries: blueberry maggot, cherry and cranberry fruitworms, and Japanese Beetles.

Cranberries: cut worms, fireworms, fruitworms, Japanese beetle and leafhoppers.

Beans: (Green, Lima, Snap beans, cowpeas and blackeyed peas) Mexican bean beetle, bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar and western bean cutworm, armyworms, corn earworm, stink bugs and tarnish plant bug.

Cucumber, Melons, Pumpkin and Squash: pickleworm, melonworm, cucumber beetles, flea beetles, leafhoppers, and squash bugs. May cause some leaf injury during prolonged high humidity. Do not use on watermelons in Florida.

Tomato, Eggplant and Pepper: Colorado potato beetle, flea beetles and leafhoppers, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

Okra: corn earworm and stink bugs. Repeat at 5-7 day intervals.

Carrots: flea beetles, harlequin bug and leafhoppers, six-spotted leafhopper (aster yellows vector), armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs and stink bugs.

Cabbage, Broccoli, Brussels Sprouts and Cauliflower: flea beetles, harlequin bug, armyworms, cabbage caterpillars and corn earworm.

Dogs and Cats: Ticks and fleas.

Dust liberally over entire animal, avoiding eyes, rub thoroughly into skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Also, dust liberally in and around sleeping quarters, kennels and dog houses. Be sure to force dust into cracks and crevices. Repeat as necessary. Avoid treatment of kittens and puppies less than 4 weeks old.

Lawn Insects: Dust liberally using 1 lb. per 275 sq. ft. for control of ants, chinch bug, earwigs, fall armyworm, fleas, leafhopper, mosquitoes, sod webworms, lawn moths, millipedes. For best results mow lawn and apply after watering or rain. Repeat in 2-3 weeks if necessary.