

Plums, Prunes and Cherries: black cherry aphid, mealy plum aphid, cherry fruit fly, cherry fruitworm, eye-spotted bud moth, fruit tree leaf roller, red-banded leaf roller, Japanese beetle, lesser peach tree borer, peach twig borer, plum curculio, prune leafhopper, brown soft scale, Forbes scale, leconium scales, oystershell scale and San Jose scale.

Peaches, Apricots and Nectarines: codling moth, cucumber beetles, European earwig, Japanese beetle, Pandemis moth, June beetles, lesser peach tree borer, orange tortrix, oriental fruit moth, peach twig borer, periodical cicada, plum curculio, fruit tree leaf roller, red-banded leaf roller, Platynota flavendana, tussock moth, Lecanium scales, San Jose scale and olive scale.

Apples and Pears: Apple sucker, green apple aphid, wooly apple aphid, bagworm, California pear-slug, codling moth, apple rust mite, pear leaf blister mite, pear rust mite, pear psylla, tentiform leaf miners, San Jose scale. Do not use until 30 days after full bloom

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust or spray mist. Avoid prolonged or repeated contact with skin. Avoid contamination of foodstuffs. Avoid contamination of feed, feeding troughs and watering receptables. Do not use after oil sprays. Wash vegetables before use.

Wash thoroughly after using.

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate) 5%
Sulphur 80%

INERT INGREDIENTS 15%

* Trademark of Union Carbide Corporation
SEVIN U.S. Patent No. 2,903,478

WONDER CHEMICAL CO., INC.



VEGETABLE DUST

Contains SEVIN

INSECT CONTROL FOR MANY HOME GARDEN INSECTS

HARVEST LIMITATIONS

No Limitations: on Beans, Blueberries, Carrots, Cucumbers, Eggplants, Grapes, Melons, Peppers, Potatoes, Pumpkins, Squash, Tomatoes, and Okra.

Allow 1 Day between spraying and harvest of: Apples, Cherries, Cranberries, Peaches, Pears, Plums, and Strawberries.

Allow 3 days between spraying and harvest of: Cabbage, Cauliflower, Kohlrabi, Lettuce, Brussels Sprouts, Broccoli, Nectarines and Apricots.

Allow 5 days between spraying and harvest of: Citrus Fruits.

W A R N I N G . . .
KEEP OUT OF REACH OF CHILDREN

DIRECTIONS: Read entire label.

Apply Vegetable Dust lightly to top and bottom of leaves when insects first appear. A light application and good coverage will give good control. Apply at weekly intervals until pests are controlled.

Beans: (Green beans, lima/snap, cowpeas, and black-eyed peas), bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvet bean caterpillar, western bean cutworm, armyworm, corn earworm, stink bugs and tarnished plant bug.

Potato, Tomato, Eggplant and Pepper: Colorado potato beetle, flea beetle, leafhopper, European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs.

Okra: corn earworm and stink bugs

Cabbage, Broccoli, Brussels Sprouts, Cauliflower and Kohlrabi: flea beetles, harlequin bug, army worms, cabbage caterpillars and corn earworm.

Lettuce and Carrots: flea beetles, harlequin bug and leafhoppers, six spotted leafhopper (aster yellows vector), armyworms, cabbage caterpillars, corn earworm, spittlebugs, tarnished plant bugs and stink bugs

Cucumber, Melons, Pumpkin and Squash: cucumber worm, melonworm, cucumber beetles, flea beetles, leafhoppers and squash bugs

Strawberries: meadow spittlebug, strawberry leaf roller and strawberry weevil

Grapes: European fruit lecanium, grape leaf folder, grape leafhopper, and grape leaf skeletonizer

Plums, Prunes and Cherries. Insects which damage plum, apricot, cherry, and peach trees include the following: eye spotted, bark, and scale insects; and rosette, red-banded, leaf-miner, spotted leaf-miner, lesser peach tree borer, spruce twig borer, plum curculion, prune leaf hopper, tree scale, leaf miner, European spruce sawfly, European spruce sawfly, scale and San Jose scale.

Peaches, Apricots and Nectarines. Insects which damage peach, apricot, and nectarine trees include the following: European sawfly, plum curculion, beetle, Prune, plum, and cherry beetles, lesser peach tree borer, orange tortrix, and the peach twig borer. Periodic applications of this product to the leaves of the trees will control these insects. San Jose scale and the European spruce sawfly are also controlled by this product.

Apples and Pears. Apples and pears are damaged by the apple aphid, sawfly, leaf miner, pear slug, codling moth, apple and pear leaf miner, and the pear leaf-miner. Pear leaf-miner is controlled by this product. San Jose scale and the European spruce sawfly are also controlled by this product.

CAUTION: Do not spray on windward side of trees. Do not spray on wet foliage. Do not spray on plants which are stressed or have been damaged by frost. Do not spray on plants which are in bloom. Do not spray on plants which are in fruit. Do not spray on plants which are in leaf fall. Do not spray on plants which are in winter dormancy.

Do not spray on plants which are in bloom.

ACTIVE INGREDIENTS:
 1. 1,1,1-Trichloro-2,2,2-trifluoroethane
 2. 2,4-Dichlorophenoxyacetic acid

INERT INGREDIENTS:



Contains "SEVIN"

INSECT CONTROL FOR MANY HOME GARDEN INSECTS

HARVEST LIMITATIONS

No Limitations: on Beans, Blueberries, Carrots, Cucumbers, Eggplants, Grapes, Melons, Peppers, Potatoes, Pumpkins, Squash, Tomatoes, and Okra.

Allow 1 Day between spraying and harvest of: Apples, Cherries, Cranberries, Peaches, Pears, Plums, and Strawberries.

Allow 3 days between spraying and harvest of: Cabbage, Cauliflower, Kohlrabi, Lettuce, Brussels Sprouts, Broccoli, Nectarines and Apricots.

Allow 5 days between spraying and harvest of: Citrus Fruits.

WARNING
KEEP OUT OF REACH OF CHILDREN

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Cabbage, Broccoli, Brussels sprouts, Cauliflower, and Kohlrabi. Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.

Do not spray on plants which are in bloom.