



DIRECTIONS FOR USE

Dilution: 2 1/2 tablespoons Rose Spray per gallon of water. Regular spraying with Rose Spray at 7 to 10 day intervals is recommended. During periods favoring powdery mildew infestation, it may be necessary to spray at weekly intervals. Do not spray if temperatures above 95 degrees F.

ROSES: Kills aphids, thrips, red spider, white flies, Japanese and May beetles, cucumber beetles, rose chafers, and controls powdery mildew. Early insect and powdery mildew control on roses is necessary as an aid to producing healthy, fast growing, and early blooming roses.

SHRUBS: To control such insects as white flies, lace bugs, spider mites, birch and holly leaf miners, leafhoppers, May beetles, Japanese beetles, webworms, tent caterpillars, powdery mildew on crape myrtle and photinia, spray thoroughly.

FLOWERS: To control aphids, spider mites, leaf miners, leafhoppers, armyworms, cutworms, thrips as well as powdery mildew, spray when insects or disease appear. Repeat as necessary.

WARNING: Do not use on food crops. Do not use on humans or pets. Contact with skin can cause toxic symptoms. Do not breathe spray mist or allow skin contact. In case of spillage on skin, wash immediately with soap and water. Poisoning may occur as a result of exposure. Avoid contamination of feed and foodstuffs. May be fatal if swallowed, inhaled, or absorbed through skin. Keep out of reach of children. Keep children and pets away from treated areas and do not handle plants until dry.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be harmed. Keep out of lakes, ponds, and streams. Do not apply to any area not specified on the label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

WONDER CHEMICAL COMPANY

San Antonio, Texas

E. P. A. Reg. No. 7393-41

- With Systemic Insecticide for Aphids, Thrips, Spider Mites
- Karathane for Powdery Mildew
- Systemic Insecticide is Absorbed into Leaves and Translocated to all Parts of Plant
- Can be Used on Flowers and Shrubs for Insect and Powdery Mildew Control



ROSE SPRAY

WITH SYSTEMIC INSECTICIDE

ACTIVE INGREDIENTS:

*Methoxychlor, technical	12.50%
*S [2 (Ethylsulfanyl)ethyl] O, O dimethyl phosphorothioate	6.25%
*2 (1-Methylheptyl)-4, 6-dinitrophenyl crotonate	1.40%
And other nitrophenols and derivatives, chiefly,	
1-methylheptyl-4, 6-dinitrophenol	0.15%
Aromatic Petroleum Distillates	63.00%
INERT INGREDIENTS:	16.70%
Total	100.00%

- *12.50% Methoxychlor, technical, is equivalent to 11% 2, 2-bis(p-methoxy phenyl)-1, 1, 1-trichloroethane and 1.5% other isomers and reaction products.
- *Meta-Systox-R reg. trademark Forbenfabriken Bayer, A.G., Chemagro Corp. licensee U.S. Pat. No. 2,791,699
- *Karathane reg. trademark Rohm & Haas

WARNING: KEEP OUT OF REACH OF CHILDREN
See other warnings on back of label.

1914

1914

1914

1914