

74101-10

P11-16

12/17/81

133

(FRONT PANEL)

ferti-lome (R)

MALATHION
GARDEN SPRAY

Contains Spreader-Sticker

Controls Aphids, Thrips, Bagworms, Houseflies, Fleas and many other insects.

Can be used on Ornamentals, Shrubs, Vegetables, Certain Trees and other plants.

2 Teaspoonsful Make 1 Gallon of Spray

ACTIVE INGREDIENTS:

Malathion (0,0-diethyl dithiophosphate of diethyl Mercaptosuccinate)55.0%

INERT INGREDIENTS:.....45.0%

(Inert Ingredients include the following.)

Octyl phenoxy polyethoxy ethanol.....0.5%

Modified phthalic glycerol alkyd resins.....0.5%

Aromatic petroleum hydrocarbons.....34.0%


Total.....-.....100.0%

CAUTION

Keep Out Of Reach of Children

See Back Panel for Additional Cautions.

Not Reviewed. Registrant claims to be in accordance with EPA's Malathion Pesticide Product Guidance for Phase 2 (see page, Page 2.12, Active to Inert Change in Status.



1. What is this for use for?

This package of ferti-lome Malathion Spray controls aphids, thrips, spider mites, lace bug, Japanese beetle adult, tent caterpillar, soft scale, bagworms, houseflies and fleas.

2. How do you use it?

Mix ferti-lome Malathion Spray (all the way through) with water by agitation; agitate the mixture while spraying. If allowed to stand, agitate before use.

ON FRUIT TREES: For control of codling moth, the curculio, apple maggot, and other fruit-eating insects, apply 2 to 3 pints of ferti-lome Malathion per 100 gallons of water per acre. Controls green apple aphid, red-banded leaf miner, European spruce sawfly, cover mite, apple tree scale, apple tree louse, apple tree cicada, apple tree psyllid, apple tree thrips, apple tree mite, and apple tree scale. Do not apply to fruit trees within 3 weeks of harvest, or to peaches within 7 days of harvest. This spray application may cause injury to McIntosh and Cortland apples within 4 weeks of harvest, or to Northeast or Rose pears under certain conditions. Consult State Extension Service Specialists for additional use on fruit trees.

ON FRUIT: For field application, apply 2 to 3 pints per 100 gallons of water per acre. Spray for fruit trees should be applied at a rate of 1000 to 1200 gallons per acre on apples and pears and 500 to 800 gallons per acre on peaches.

To control spider mites and aphids on tomatoes, beans, corn cobs, potatoes, use 1 1/2 teaspoonful of Malathion per gallon of water. Use same to get complete coverage. Do not apply within 7 days of harvest. For field application apply 1 1/2 pints ferti-lome Malathion per acre in sufficient water to give thorough coverage.

On ornamentals, to control lace bug, weevil, codling moth, aphids, mealybugs, spider mites, whiteflies, fourlined leaf bug, Japanese beetle adult, potato leafhopper, tarnished plant bug, thrips, red-banded leaf miner, and curly scale, mix 1 1/2 teaspoonful per gallon of water and spray thoroughly. Do not use on certain species of crabapple, and on certain species of deciduous, holly and yew.

On ornamentals to control birch leaf miner, boxwood leaf miner, tent caterpillar, azalea scale, pine leaf scale, and soft scale, mix 1 1/2 pints per gallon of water and spray thoroughly.

ON HOUSES: For control of house flies, mix 1 1/2 pints of ferti-lome Malathion per 100 gallons of water and spray thoroughly. Use same to get complete coverage. Do not use on certain species of crabapple, and on certain species of deciduous, holly and yew.

For control of fleas on cats, dogs, mice, rats, and under houses, mix 1 1/2 pints of ferti-lome Malathion per 100 gallons of water and apply at rate of one gallon of mixture per 1000 square feet, use garden sprayer.

Back label can't

3. Here are the results you may expect:

Ferti-loke Malathion, used according to directions, allows you to control insect pests in the garden, on shrubs, on flowers and on fruit trees. This new effective spray can be a great help to you in solving many insect control problems.

Directions: Normal application rate is 1/2 ounce per gallon of water. Avoid contact with skin, eyes or clothing. Do not use on edible plants. Do not use on plants that are stressed or in bloom. Do not use on plants that are susceptible to phytotoxicity. Do not use on plants that are susceptible to malathion. Do not use on plants that are susceptible to malathion.

Caution: Do not use on plants that are stressed or in bloom. Do not use on plants that are susceptible to phytotoxicity. Do not use on plants that are susceptible to malathion. Do not use on plants that are susceptible to malathion.

NON-WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

7901-1

The Ferti-loke Co.

Houston, Texas

1968

