Customer's Signature

ACCEPTE


NOV I 1 1971

UNDER THE FEDERAL INSECT. FUNCTORS AND PERSONS FOR ECONOMIC FOR THE REST ED UNDER NO 10.05. I TO ATTACHED COME.

1401-14

FOR BEST RESULTS HOLD AT 45° ANGLE TO GROUND







ONDERY MILDEW SFIDER: MITES! THRIPS ANDICERTAIN OTHER अग्राताणाः तक्ताक તારે લગ્સ્સ AGON STREET NEW TO WAR IN COMMENT OF THE PROPERTY OF THE PROPE . XI IV.

्योतः विदयस्योग्यास्योग्यास्यात्रस्याः Applitu.

नरावरा नर्तितवन।वराहरू नेवा

14. 6 iliteration to

 $\mathcal{J}_{1}(x_{1})$.

GAUMON

CHANGE CONTRACT OF THE CONTRAC त्यः वातः भाषा स्वाभावाताताम् व्यापाताह

ROSE DUST AND SPRAY

Serves as a hundy Garden Dust Gun

For refills, purchase the economical 2 lb. can.

Remove safety cap, point dust gun toward plant . . and squeeze.

WHAT IS THIS PACKAGE FOR?

package of ferti-lome ROSE DUST AND SPRAY is to be for control of insects and diseases in roses, flowers and snrubs. It may be used as a dust or spray.

INSECTS CONTROLLED: Thrips, Aphids, Bagworms, Grass:
hoppers, Crickets, May beetles; Chimbing Cutworms, Leaf beetles,
Leaf Rollers, Lacebugs, Japanese beetles, Rose chafer, Rose
wacyrs, Spider miles, Red Spider, Rose Leafhopper, Rose slugs;
also controls Azalea scale, Scuriy scale, Euonymus scale. Softbrown scale, Boxwood Leaf-miners and many others.

DISEASES CONTROLLED: Black Spot, Powdery Mildew, Leaf

Spot and Rust.

2. HOW DO YOU USE IT?

Dusting: Roses and other flowers - to be most effective Dusting: Roses and other flowers—to be most effective, applications should be made before insects become numerous and before diseases are apparent, Protect plants through the growing season by dusting or spraying at 7 to 10 day intervals and after heavy rains. Cover foliage thoroughly, but do not use in excess. Pay particular attention to cover underside of leaves. For best results, apply in early morning or late evening when the temperature is low and the air is still. Do not treat during periods of excessively high temperature (about 90°F) as damage to foliage may occur. Sulfur may burn foliage when temperatures are excessively high. tures are excessively high.

Spraying: Mix 8 tablespoonsful in a little water to make a thin paste. Add to 1 gallon of water and agitate during application. If used in a small sprayer, shake often for agitation. Follow dusting directions for use for timing and application.

3. HERE ARE THE RESULTS YOU MAY EXPECT.

You will find that when you follow a regular scheduled insect and disease control program, your roses and flowers will not produce more blooms, but also larger and more colorful.

CAUTION: HARMFUL IF INHALED OR SWALLOWED.

Do not breathe dust. Do not take internally, Avoid breathing of dust. Skin contact may be harmful, Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. ATROPINE IS ANTIDOTAL. For protection of honeybees, avoid unnecessary use during periods when honeybees are visiting crops. Dispose of container promptly when empty. KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

This product is toxic to fish. Keep out of lakes, streams or ponds. Apply this product only as specified on this label. NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or randling of this material not in strict accordance with directions given herewith.

EPA . Reg. No. 7401-14

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS

NET CONTENTS 10 OZ.

"Sevin" Reg. Trademark Union Carbide Corporation "Keithane" Reg. Trademark Rohm & Haas Company