



Makers of fine lawn & garden and farm chemicals / Natural Guard Brand / Fermers Cooperative Brand / ferti-tome Brand / AMERICAN Brand / Ht-Yield Brand

Feb. 22, 1993

Joanne I. Miller, Product Manager 23 U.S. EPA OPP Registration Division (H7505C) 401 M Street SW Washington, D.C. 20460

I, Being an authorized representative of Voluntary Purchasing Groups, Inc. certify that all containers fo ferti-lome Lawn Weed Killer Granules, 7701-334, produced by June 15,1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said products sold or distributed by this company by jan. 1,1995 will bear revised labeling in accordance with the revised labeling required for Task Force Technical and manufacturing-use products.

Sincerely,

Raymond H. Smith

Raymond H. Smith Product Manager

(Upper Back Panel) p.3

Net Wt. 30 Lbs.

ferti-lome (R)

## LAWN WEED KILLER GRANULES

FOR CONTROLLING WEEDS IN COMMON BERMUDA, BAHIAGRASS, ZOYSIA, KENTUCKY BLUE, MERION BLUE, RED TOP, CREEPING RED FESCUE, ST. AUGUSTINE GRASS AND RYE GRASS LAWNS

EPA Est. 7401-TX-1

EPA Reg. No. 7401-33

VOLUNTARY PURCHASING GROUPS, INC. Bonham, Texas 75418

(Lower Back Panel)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION: It is a violation of Federal Law to use this product in a manner inconsistent with it's labeling.

1. WHAT IS THIS PACKAGE FOR?
This package of ferti-lome LAWN WEED KILLER GRANULES is an easy to use granular form for controlling broad leaf and vining type weeds. It can be used to control weeds in lawns, golf courses, and similar areas. The following is a partial list of weeds controlled:

| Landelion      | Henbit              | Mouseear Chickweed |
|----------------|---------------------|--------------------|
| Clovers        | Oxalis              | Spotted Spurge     |
| Chicory        | Mustard             | Stitch Wort ::     |
| Docks          | Pennywart           | Wild Aster         |
| Dog Fennel     | Peppergrass         | Wild Carrot        |
| English Daisy  | Pigweed             | Wild Garlic        |
| Frenchweed     | Plantain (Buckhorn, | Wild Onion         |
| Erect Knotweed | Broadleaf, Rugel's) | Wild Radish        |
| Ground-Ivy     | Poison Ivy          | Wood Sorrel        |
| Hawkweed       | Poison Oak          | Yarrow             |
| Healall        | Prostrate Knotweed  | Puncturevine       |
| Lambsquarter   | Purslane            | Milkweed           |
| Mallow         | Ragweed             | Bull Thistle       |
| Morning Glory  | Sheep Sorrel        | Shepherd's Purse   |

# (Lower Back Panel)

2. HOW DO YOU USE IT?

Apply uniformly to area to be treated. For lawn and similar areas the spreader setting below as a guide. Apply at rate of 6 lbs. per 1000 Sq. Ft.

Spreader Setting\* Spreader Setting\*
Spreader Coverage Once over Twice over
20" Deluxe 5000 Sq.Ft. 6 4
Cyclone 5000 Sq.Ft. 3 2

Best results are obtained by applying when weeds are wet from dew, sprinkling, or rain. Control is slower where material is watered into soil and absorbed into system through roots. However, it is effective when absorbed through roots. For St; Augustine grass apply one 30 lb. bag on 20,000 Sq. Ft. Repeat application in 10-14 days if necessary. Heavily infested lawns may require repeat application to kill missed weeds. Best results are obtained when weeds are young and actively growing. Application in early Spring and Fall when moisture and temperature make weeds grow rapidly is a good time to apply. Watering and fertilizing schedules should be maintained before and after application. Do not apply during periods of drought. Do not apply within the dripline of desirable trees and shrubs.

If lawn begins to brown temporarily, water heavily. Do not use on Centipede grass, Dichondra, nor on lawns or turf where desirable bentgrass or clovers are present. Do not use on newly seeded lawns or turf.

\*The settings for spreaders can only be taken as approximate. Our products have been thoroughly tested on new spreaders only. Age and condition of spreader, speed of pushing, roughness of lawn, all affect application. Operator should carefully observe rate of application at beginning of job and make corrective adjustments of spreader opening if necessary.

Application can be made on dry lawns or similar areas where weeds are to be controlled. When such an application is made, thorough watering should follow. Do not "puddle"

3. HERE ARE THE RESULTS YOU MAY EXPECT! ferti-lome LAWN WEED KILLER GRANULES used according to directions gives you economical and effective control of weeds in your lawn thus letting your lawn grasses grow vigorously to provide you with a beautiful weed free lawn.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.



## (Upper Front Panel)

# controls certain broadleaf and vining type weeds

Covers up to 5,000 Sq. ft.

- \* NEW SYSTEMIC ACTION
- \* ATTACKS THROUGH LEAVES
- \* ATTACKS THROUGH ROOTS

(Lower Front Panel)

# LAWN WEED KILLER GRANULES

FOR CONTROLLING WEEDS IN COMMON BERMUDA, BAHIAGRASS, ZOYSIA, KENTUCKY BLUE, MERION BLUE, RED TOP, CREEPING RED FESCUE, AND ST. AUGUSTINE GRASS AND RYEGRASS LAWN GRASSES.

| ACTIVE INGREDIENTS: Dimethylamine Salt of Dicamba (3,6-dichloro-o-                  |
|---|
| anisic acid)*   |
| acid  |
| INERT INGREDIENTS: 33.07%   |
| Total130.00%  |
| *Equivalent to .19% 3,6-dichloro-o-anisic acid (dicamba). +Equivalent to 2,4-D Acid |

KEEP OUT OF REACH OF CHILDREN (12 point type)
CAUTION (18 point type)



PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

5 of 5

CAUTION: May be harmful or fatal if swallowed. Avoid contact with skin, eyes and clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

#### STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED- Call a physician or Poison Control Center immediately.

if possible, vomiting should be induced under medical
supervision. Drink one or two glasses of water and induce
Vomiting by touching back of throat with finger. Do not
induce vomiting or give anything by mouth to an unconscious
person.

IF INHALED - Remove victim to fresh air. Apply artificial respiration if indicated.

IN ON SKIN - Remove contaminated colthing and wash affected areas with soap and water.

IF IN EYES - Flush eyes with plenty of water. Call a physician immediatel

#### **ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drifts or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and pothles). Do not contaminate water when disposing of equipment washwaters. Most case of groundwater contamination involving phenoxy herbicides such as 2,4-D have associated with mixing/loading and disposal sites. Caution should be exercisedawhen handling 2,4-D pesticidesat such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pai to contain spills will help prevent groundwater contamination.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

VOLUNTARY PURCHASINGGROUPS, INC.

BONHAM, TEXAS 75418

NET WEIGHT \_\_\_\_ LBS.

