

pm23

7401-37

145



Makers of fine lawn & garden and farm chemicals / Natural Guard Brand / Farmers Cooperative Brand / ferti-lome Brand / AMERICAN Brand / Hi-Yield Brand

Feb. 22, 1993

Joanne I, Miller, Product Manager 23  
U.S. EPA  
OPP  
Registration Division (H7505C)  
401 M Street SW  
Washington, D.C. 20460

I, Being an authorized representative of Voluntary Purchasing Groups, Inc., certify that all containers of ferti-lome Weed & Feed for controlling Weeds, ~~7401-37~~, produced by june 15, 1994 will bear revised labeling in accordance with the revised labeling required for Task Force technical and manufacturing-use products. I further certify that all containers of said products sold or distributed by this company by Jan. 1, 1995 will bear revised labeling in accordance with the revised labeling required for Task Force Technical and manufacturing-use products.

Sincerely,

A handwritten signature in cursive script that reads "Raymond H. Smith".

Raymond H. Smith  
Product Manager

(Upper Front Panel)

controls weeds . . . and feeds your lawn

Coverage  
3,000 Sq.ft.

- \* NEW SYSTEMIC ACTION
- \* ATTACKS THROUGH LEAVES
- \* ATTACKS THROUGH ROOTS

(Lower Front Panel)

WEED & FEED

FOR CONTROLLING WEEDS  
IN

COMMON BERMUDA, ST. AUGUSTINE, BAHIA, ZOYSIA, KENTUCKY BLUE,  
MERION BLUE, RED TOP, CREEPING RED FESCUE, AND RYE GRASS LAWN  
GRASSES

ACTIVE INGREDIENTS:

Dimethylamine Salt of Dicamba (3,6-dichloro- o-anisic acid)* -----	.11%
+Dimethylamine Salt of 2,4-Dichlorophenoxyacetic acid -----	.31%
INERT INGREDIENTS: -----	99.58%
Total -----	100.00%

\*Equivalent to .09% 3,6-dichloro-o-anisic acid (dicamba)

+Equivalent to 2,4-D acid ----- .26%

KEEP OUT OF REACH OF CHILDREN (12 point type)  
CAUTION (18 point type)

PRECAUTIONARY STATEMENTS

3 of 5

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** May be harmful or fatal if swallowed. Avoid contact with skin, eyes and clothing. When using this product, wear long-sleeved shirt, long pants, socks, shoes and chemical resistant gloves. After using this product, rinse gloves before removing, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water. Remove saturated clothing as soon as possible and shower. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment areas until spray has dried or dust has settled.

**STATEMENT OF PRACTICAL TREATMENT**

- IF SWALLOWED-** Call a physician or Poison Control Center immediately. if possible, vomiting should be induced under medical supervision. Drink one or two glasses of water and induce Vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.
- IF INHALED** - Remove victim to fresh air. Apply artificial respiration if indicated.
- IF ON SKIN** - Remove contaminated clothing and wash affected areas with soap and water.
- IF IN EYES** - Flush eyes with plenty of water. Call a physician immediately.

**ENVIRONMENTAL HAZARDS**

This product is toxic to aquatic invertebrates. Drifts or runoff may adversely affect aquatic invertebrates and nontarget plants. Do not apply directly to water or wetlands (swamps, bogs, marshes and potholes). Do not contaminate water when disposing of equipment washwaters. Most cases of groundwater contamination involving phenoxy herbicides such as 2,4-D have associated with mixing/loading and disposal sites. Caution should be exercised when handling 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing or transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

Buyer assumes all risks of use, storage or handling of this product not in strict accordance with directions given herewith.

**VOLUNTARY PURCHASING GROUPS, INC.**

**BONHAM, TEXAS 75418**

NET WEIGHT \_\_\_\_\_ LBS.

**BEST AVAILABLE COPY**

(Upper Back Panel)  
NET WT. 40 LBS.

475

(Upper Back Panel)

Net Wt. 40 Lbs.

ferti-lome (R)

WEED & FEED  
FOR CONTROLLING WEEDS IN  
COMMON BERMUDA, ST. AUGUSTINE, BAHIA, ZOYSIA, KENTUCKY BLUE,  
MERION BLUE, RED TOP, CREEPING RED FESCUE, AND RYE GRASS LAWN  
GRASSES.

11-5-6

GUARANTEED ANALYSIS

Total Nitrogen (N) -----	11.00%
11.00% Ammoniacal Nitrogen	
Available Phosphoric Acid (P <sub>2</sub> O <sub>5</sub> ) -----	5.00%
*Soluble Potash (K <sub>2</sub> O) -----	6.00%

Derived from - Primary Plant Foods: Sulphate of Ammonia,  
Ammoniated Superphosphate, Muriate of Potash.

\*In states requiring, reads Water Soluble (K<sub>2</sub>O).

EPA Est. 7401-TX-1

EPA Reg. No. 7401-37

VOLUNTARY PURCHASING GROUPS, INC.  
Bonham, Texas 75418

(Lower Back Panel)

DIRECTIONS FOR USE

GENERAL CLASSIFICATION: It is a violation of Federal Law to  
use this product in a manner inconsistent with it's labeling.

Dear Folks:

We recommend this package of ferti-lome WEED & FEED FOR CON-  
TROLLING WEEDS because we have found it most effective. I  
think you will find the following information complete and  
easy to follow.

1. WHAT IS THIS PACKAGE FOR?

ferti-lome WEED & FEED FOR CONTROLLING WEEDS is for applying  
to your lawn when you want to give it a good feeding and at  
the same time kill weeds.

BEST AVAILABLE COPY

5 of 5

(Lower Back Panel)

p. 3

2. HOW DO YOU USE IT?

For best results apply when temperature is 65°F or above. Apply one 40 lb bag per 3000 square feet of lawn area. Apply uniformly using fertilizer spreader. Apply when grass is dry and water in thoroughly. See Chart below for proper ferti-lome spreader setting to apply this product. Apply when weeds have appeared and before they reach maturity. Shut off spreader when stopping or turning around.

SPREADER	ONE BAG COVERS 3,000 sq. ft.	
	SPREADER SETTING*	
ferti-lome Cyclone	4 1/2	to build good turf
ferti-lome Deluxe	8	feed with ferti-lome
Jackson	7	monthly during
Lawn Beauty	6	growing season
Scotts	6 1/2	
Sears	8	
Central	8	

\*These settings can be used only as a guide, they are not absolute. The speed at which the applicator walks, the condition of the spreader, and the texture of the turf will alter the rate of fertilizer application.

Do not use on Centipede, Bent Grasses, Lippia or similar ground covers. Do not use on Dichondra, Weed killing action is most effective when weeds are growing rapidly. This is normally during time of warm weather, adequate moisture and plant food. Treatment during dry or cool weather usually reduces killing action.

New lawns should be well established and mowed at least twice before treatment. This product controls clover. Do not use on lawns where clover is desired. Clean spreader thoroughly after using.

For use on ST. Augustine grass apply 1-40 lb. bag per 9,000 sq. ft. of lawn area. Repeat application if necessary in 10-14 days.

3. HERE ARE THE RESULTS YOU CAN EXPECT!

ferti-lome WEED & FEED FOR CONTROLLING WEEDS applied according to directions will give you control of dandelions, henbit, chickweed, buckhorn, plantain, clovers, smart weed, wild carrot, mustard, Ground Ivy, English Daisy, Filaree and other similar weeds and the same time measurably improve your lawn due to the application of a complete plant food.

Sincerely,

the ferti-lome people

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

BEST AVAILABLE COPY