

7401-38

09/22/1971

1/2

NEW!

(Front Panel)

ferti-lome

Liquid

SEVIN* GARDEN SPRAY

* CAN BE USED ON ORNAMENTALS, LAWNS, FRUIT TREES, SHADE TREES, VEGETABLES AND CITRUS TREES.

* 4 TEASPOONFULS MAKE 1 GALLON OF SPRAY FOR CERTAIN USES.

ACTIVE INGREDIENT:

Carbaryl, (1-naphthyl N-methylcarbamate) 23.7%

INERT INGREDIENTS: 76.3%

NET CONTENTS 1 PINT LIQUID

CAUTION (18 Point)

KEEP OUT OF REACH OF CHILDREN (12 Point)

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS (12 Point)

(Side Panel)


1. WHAT IS THIS PACKAGE FOR:

ferti-lome Liquid SEVIN GARDEN SPRAY is an easy to use insecticide for use on Ornamentals, Lawns, Fruit Trees, Shade Trees, Vegetables and Citrus Trees.

2. HOW DO YOU USE IT?

ROSES, ORNAMENTALS, FLOWERS, SHRUBS, SHADE TREES: Shake well before using or sampling. Mix 4 teaspoonfuls per gallon of finished spray desired. For control of Japanese Beetle, Aphids, Bagworms, Elm Leaf Beetle, certain Leaf Miners, Rose Slugs, Periodical Cicada (17 year locust), Scale insects, Tent Caterpillars, Blister Beetles, Leaf Rollers, and exposed Thrips. Start spraying when insects or damage appear. Spray thoroughly covering both sides of leaves. Repeat application in 7-10 days as necessary to maintain control.

VEGETABLES: Mix 4 teaspoonfuls per gallon of finished spray desired. For control of Pickleworm, Fall Army Worms, Corn Borer, Harlequin Bug, Fruitworms, Melon Worms, Horn Worms on tomatoes, eggplant, peppers, cucumbers and summer squash. Do not apply within 1 day of harvest.



2/2

(Side Panel)

Page 2.

FRUIT & CITRUS TREES: Mix 4 teaspoonfuls per gallon of finished spray desired. For control of Codling Moth, Cucumber Beetles, Fruit Tree Leaf Roller, Lesser Peach Tree Borer, Plum Curculio, Red-Banded Leaf Roller, Apple Maggot, Bagworm, Citrus Cutworm, Orange Tortrix on Apple, Pear, Citrus, Peach, Apricot and Nectarine. On Fruit Trees do not apply with 5 days of harvest.

LAWNS: Mix 1 pint in 15 gallons of water and apply to 500 sq. ft. of lawn area. For best results apply to mowed lawn. For control of Chinch bugs, Sod Webworms, Fall Army Worms, Fleas and Ants. Repeat in 2-3 weeks if necessary. Do not allow children, pets or domestic animals on treated lawn until the insecticide is washed off the grass into the soil by sprinkling or rainfall and grass is completely dry.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

ferti-lome Liquid SEVIN GARDEN SPRAY used according to directions, controls certain insects on plants, trees and other areas.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of spray. Do not take internally. Skin contact may be harmful. Avoid contact. Wash hands and face before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Do not reuse container for any purpose. Destroy empty container by breaking and burying in a safe place.

Food utensils such as teaspoons and tablespoons should not be used for food purposes after use with pesticides.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATER RECEPTACLES.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

BEE CAUTION: For protection of honey bees avoid use, if possible, during periods when honey bees are visiting the crop. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until 1 week after application.

NOTE: Product freezing temperature 32° F. Satisfactory utility is restored after thawing and mixing of product. Repeated 0-10° freezing-thawing may impair utility.

* Trademark of Union Carbide Corporation for the active ingredient 1-naphthyl N-methylcarbamate (carbaryl).

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS USDA REG. NO. 7401-00