

6/20/97

PM 19

7401-43

Pg 175

Front Panel

ferti-lome
INSECT KILLER
(Granular)

Contains SEVIN* Insecticide

\$5.95
Covers up to
5000 Sq. Ft.

FOR THE CONTROL OF CHINCH BUGS, SOD WEBWORMS, AND CERTAIN OTHER INSECTS

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate).....	3.34%
INERT INGREDIENTS:.....	96.66%
Total.....	100.00%

*SEVIN is the registered trademark of Rhone Poulenc for carbaryl insecticide.

CAUTION ---18 point type

Keep Out Of Reach Of Children---14 point type
See Side Panel For Additional Cautions---14 point type

the ferti-lome people

A DIVISION OF

VOLUNTARY PURCHASING GROUPS, INC.

Bonham, Texas

Net. Wt. 30 Lbs.

ACCEPTED
JUN 20 1997
 Under the Federal Insecticide,
 Fungicide, and Rodenticide Act,
 as amended, for the pesticide
 registered under
 EPA Reg. No. 7401-43

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. READ ALL DIRECTIONS FOR USE AND PRECAUTIONARY STATEMENTS BEFORE USING.

KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT INTO FOOD OR DRINK CONTAINERS.

USER SAFETY REQUIREMENTS

- * Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- * Change clothing as soon as possible after use.
- * Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- * Do not allow children or pets to contact treated areas until dust has settled.

HOME LAWNS, RECREATIONAL, AND ORNAMENTAL TURF AREAS

For control of turfgrass pests, apply granules uniformly to the turf grass. For optimum control, follow specific directions for proper watering or irrigation after application. Do not allow public use of treated areaduring application.

INSECT	POUNDS OF SEVIN(R) GRANULES PER 1,000 SQ.FT.	SPECIFIC DIRECTION.
Ants, Armuworms, Centipedes, Chiggers, Chinchbugs, Crickets, Cutworms, Darking Ground beetles, Earwigs, Esses Skipper, European Chafer, Fall armyworms, Liery skipper, Fleas, Grasshoppers, Green June Bugs, Green June beetle grubs, Lawn Moths, (sod Webworms), Leafhoppers, Lucerne th larvae, Millipedes, Sowbugs, Spittlebugs, Springtails, Ticks, Yellowstriped armyworms.	6 to 7½	For general insect control apply granules uniformly when grass is dry. For most effective control, lawn should be watered lightly after application to remove the insecticide from the granule. Repeat applications may be made at weekly intervals to control new infestations.
White grubs (such as Japanese beetle grubs, Chafer beetle grubs, Phyllophage spp grubs, and Annual white grubs), Bluegrass billbug larvae, European crane-fly larvae.	7½	For white grub control applications should be made when grubs are feeding near the soil surface usually during the late march through May or July to early September, or as recommended by local agricultural extension agents. For most effective contro turf should be watered thoroughly after treatment to remove insecticide from

granule. Repeat applications may be made once per week.

NUISANCE PEST CONTROL IN AND AROUND FLOWER BEDS AND ORNAMENTAL PLANTINGS AND AROUND HOMES.

For control of ants, armyworms, ticks, cutworms, centipedes, crickets, earwigs, firebrats and silverfish, fleas, millipedes, sowbugs, scorpions and spiders: apply granules uniformly to a 6 foot band around building. Apply granules at the rate of 11 ounces per 120 square feet (6X20 band) to shrub and flower beds, foundations, ornamental plantings, and lawn or soil areas immediately adjacent to boulding. Formost effective control, treated area should be lightly watered after pesticide application to remove insecticide from the granules. Pepeat applicatoin weekly to maintain control.

FRUIT AND VEGETABLE CROPS (Not for use in California)

For control of Ants, Centipedes, Crickets, Cutworms, Earwigs, Grasshoppers, Millipedes, Sowbugs (Pillbugs), Springtails, and Ticks apply 23 ounces per 1,000 ft. evenly over the soil surface. Water lightly after application to release the insecticide from the granule. Applications may be made weekly before during or after the growing season. If applying during the growing season, observe indicated pre-harvest intervals (PHI).

- 0 day PHI --- Beans, blueberries, carrots, corn, cucumber, eggplant, melons, okra, peanuts, pepper, potato, pumpkins, squash, sweet potato, tomato.
- 1 day PHI---- Asparagus, strawberries.
- 3 day PHI --- Broccoli, brussels sprouts, cabbage, cowpeas, garden beat root, head lettuce, koihirabi, parsnips, peas, radishes, salsify roots, turnip roots
- 7 day PHI --- Blackberries, dewberries, (including boysenberries and loganberries), grapes, raspberries.
- 14 day PHI --- Celery, chinese cabbage, collards, dandelion, endive, (escarole), garden beet tops, hanove salad, kale, leaf lettuce, mustard greens, parsley, salsify tops, spinach, swiss chard, turnip tops.

IMPORTED FIRE ANT CONTROL

For use as a mound treatment to control imported fire ants, apply 2.8 to 5.6 ounces over and around each fire ant mound. Irrigate mound thoroughly immediately after treatment to remove insecticide from granule and flush in down into the ground. A minimum of 1 gallon of water should be applied to each mound, but do not use more than 2 gallons of water as flooding may result. Apply water gently to avoid disturbing ants and causing migration before the chemical can be effective. For best results, apply in the early morning or late evening hours when ants are least active. Treat new mounds as they appear.

CONTROL OF TICKS WHICH TRANSMIT LYME DISEASE

For control of juvenile and adult Ixodes spp. ticks (Deer tick, Bear tick, and Black legged tick) and Amblyomma spp ticks (lone star tick), Apply granules at teh rate of 4.7 to 5.7 ounces per 1000 sq.ft. Begin applications in the late spring or early summer (May/June). Uniformly apply granules to entire lawn, perimeter wooded areas, and property boundaries including shrub beds, ornamental plantings, and wooded areas where exposure to ticks may occur. For optimum results, irrigate treated areas after application to remove insecticide from granule.

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS
CAUTION

Harmful if swallowed or absorbed through the skin. Avoid breathing dust. Do not take internally. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling this product. Do not eat, drink or use tobacco while working with this product. Wash hands and face before eating, drinking or using tobacco. Keep out of reach of children and domestic animals.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of wastes.

FIRST AID

GENERAL: Contact a physician immediately in all cases of suspected poisoning. Transport patient to a physician or hospital immediately and show a copy of this label to the physician. If poisoning is suspected in animals, contact a veterinarian.

IF SWALLOWED: Never give anything by mouth to an unconscious person. If conscious and not convulsing, drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger.

IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists.

IF ON SKIN: Wash thoroughly with soap and water.

IF INHALED: Move from contaminated atmosphere and call a physician.

NOTE TO PHYSICIAN: Carbaryl is a carbamate insecticide, which is cholinesterase inhibitor. Atropine is antidotal. Narcotics and other sedatives should not be used. Further, drugs like 2 PAM (pyridine-2-aldoxime methiodide) are not recommended.

STORAGE: Store unused product in original container only. in cool dry area out of reach of children and animals.

CONTAINER DISPOSAL: Do not reuse empty bag. Discard bag in trash.

Sevin (R) is a registered trademark of Rhone-Poulenc.

ADD SPREADER SETTINGS TO GIVE HOMEOWNER GUIDE.

VOLUNTARY PURCHASING GROUPS, INC.
EPA REG. NO. 7401-43

BONHAM, TEXAS 75418
EPA EST. NO. 7401-TX-1

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Net Weight LBS.

NON-WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-43