

ferti·lome®

PLANT FOOD

CONTAINING SEVIN®
FOR CONTROL OF SOIL INSECTS

CONTROLLED BY SEVIN® IN: CRICKETS, GRASSHOPPERS, BEETLES, CATERPILLARS, COTTAGE GEAR, FLEAS, MOSQUITOES,
AND OTHER SOIL INSECTS. ALSO CONTROLLED BY SEVIN® IN: BEETLES, CATERPILLARS, COTTAGE GEAR, FLEAS, MOSQUITOES,
AND OTHER SOIL INSECTS.

CAUTION
KEEP OUT OF REACH OF CHILDREN

DO NOT BREATHE DUST OR VAPORS. DO NOT GET IN EYES OR ON SKIN. IF CONTACTED WITH SKIN, WASH IMMEDIATELY WITH SOAP AND WATER. IF CONTACTED WITH EYES, WASH IMMEDIATELY WITH COLD WATER. IF CONTACTED WITH CLOTHING, REMOVE CLOTHING IMMEDIATELY AND WASH SEPARATELY. DO NOT SMOKE OR DRINK ALCOHOLIC BEVERAGES WHILE HANDLING. WEAR REGULAR LONG SLEEVED WORK CLOTHING.

DO NOT FEED TO LIVESTOCK OR FEED AND FOODSTUFFS.

DO NOT BURN. DO NOT INCINERATE. DO NOT BURY. DO NOT SMOKE. DO NOT DRINK ALCOHOLIC BEVERAGES WHILE HANDLING.

SEVIN® IS A MODERATELY TOXIC, REVERSIBLE, CARBAMATE PESTICIDE.

ACTIVE INGREDIENTS:
1,1-Dichloro-2,2-bis(4-chlorophenyl) ethane (D.D.T.) (N-methylcarbamate)

Net Weight:

SEVIN® is a registered trademark of Union Carbide Corporation for carbamate pesticides.

NET WT. 70 LBS.

ferti-lome®

PLANT FOOD

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Do Not Take Internally. Do Not Breathe Dust. Skin Contact May Be Harmful. Avoid Contact With Skin. Wash Hands and Face Before Eating. Wash Thoroughly After Handling. Wear Regular Long Sleeved Work Clothing. Change to Clean Clothing Daily.

AVOID CONTAMINATION OF FEED AND FOODSTUFFS.

PRECAUTION: Clean Up Spillage and Bury. Empty Bags Should Be Destroyed by Burying With Waste or by Burning. If Burned, KEEP OUT OF SMOKE.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

ACTIVE INGREDIENTS:

Carbaryl (1-Naphthyl N-methylcarbamate) 1.786%

INERT INGREDIENTS: 98.214%

Total..... 100.000%

SEVIN® is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

NET WT. 70 LBS.

fortilome

DIRECTIONS FOR USE

Sevin insecticide will not control insects if used with lime or other highly alkaline materials. Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of insect infestation and repeat at 7-10 day intervals when needed to maintain control. For disease control, begin applications before diseases appear. All dosages are given in pounds per acre. Observe use limits given for specific crops.

BEANS: Use 10 lbs. for Mexican bean beetle, the Texas bean leaf beetle, leafhoppers, armyworms, cutworms, Japanese beetle, leafhoppers, powdery mildew, and the cotton caterpillar and Western bean cutworm. Use 15-25 lbs. for army worms and tarnished plant bugs. Use 20 lbs. for corn earworm, lima bean pod borer, lygids and stink bugs. May be applied up to and including the day of harvest.

COTTON: Early season control of Cotton leafworm, cotton leafhopper, fleabeetles, striped caterpillar, thrips. Apply 10 lbs. per acre. Mid and late season control of Boll weevil, boll worm, fall armyworm, cotton leaf perforator, leaf rollers, leafhoppers, lygus bugs (light to moderate infestations in Western irrigated cotton), tarnished plant bugs, aphids. Populations will be suppressed by repeated applications of this insecticide. Apply 10-20 lbs. For pink bollworm—apply 15-20 lbs. on a 5-7 day schedule depending on the size of the cotton and degree of infestation. For salt-marsh caterpillar, apply 20 lbs. For grasshoppers—Apply 10-15 lbs. The higher dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants. Allow 7 days between last application and grazing treated fields or harvesting crop residue for use as feed or bedding for dairy or meat animals.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing dust. Avoid contact with skin. Wash thoroughly after handling. Change contaminated clothing daily. Avoid contamination of feed and feedstuffs. Do not reuse empty container. Destroy container by burying with waste or burning if possible, away from smoke or fumes.

Do not apply during bloom when bees, bees or other pollinators are active. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Avoid drift to adjoining food and forage crops.

Sulfur will cause severe fruit and leaf injury to sulfur sensitive crops. Do not use sulfur on any crop unless local use has proved safe. Sulfur may burn foliage and fruit during periods of high temperature and under certain other climatic conditions. Do not apply sulfur to sensitive crops when shade temperatures exceed or are likely to exceed 90 degrees F. The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. Consult State Agricultural Extension Service for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

50-1-46

EPA Reg. No. 7407-81-AA



ferti-lome



Sevin* Sulfur 10-40 Dust

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-Methylcarbamate)	10.0%
Sulfur	40.0%

INERT INGREDIENTS:	50.0%
Total	100.0%

Sulfur Fineness: 93% Passing No. 325 U.S.S. Sieve.

***Trade-Mark of Union Carbide Corporation for the active ingredient, Carbaryl (1-naphthyl N-Methylcarbamate).**

EPA Est. 7401-TX-1

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
See Side Panel for Additional Cautions**

**VOLUNTARY PURCHASING GROUPS, INC.
Bonham, Texas**

NET WEIGHT 50 LBS.

Hi-Yield®

Toxaphene 6 E.C.

For Insect Control on Vegetables, Field Crops and Small Grains

Contains 6 lbs. Toxaphene per Gallon

ACTIVE INGREDIENTS:

Toxaphene (Technical Chlorinated Camphene
containing 67-69% Chlorine) 59.5%
Aromatic Petroleum Derivatives Solvent 36.3%

INERT INGREDIENTS: 4.2%
Total 100.0%

EPA Est. 7401-TX-1

WARNING:

KEEP OUT OF REACH OF CHILDREN
See Side Panel For Additional Warnings

NET CONTENTS

GALLONS

GENERAL INFORMATION:

Hi-Yield TOXAPHENE 6 E.C. is an emulsifiable concentrate that must be diluted with water according to the directions below. It is intended for control of insect pests on agricultural crops. In these directions given below Hi-Yield TOXAPHENE 6 E.C. should be mixed with sufficient water to assure thorough coverage. Best control is obtained when application is made at first sign of insects or their damage appear. Applications should be repeated as necessary to maintain control.

GUIDE FOR APPROXIMATE AMOUNT OF WATER TO USE PER ACRE: For airplane and hydraulic sprayers 2 to 5 gallons. Low gallonage sprayers 5 to 10 gallons. Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

NOTE: Do not feed treated vines, forage, plants, mill trash or ensilage made from plants to poultry, livestock, dairy animals or animals being finished for slaughter.

CROP	INSECTS	RATE PER ACRE	REMARKS
BEANS Green Snap Lima	Armyworms; Bean beetles (Mexican bean, bean leaf, cucumber, wax, bean blister and flea); Bean leaf skeletonizer; Darkling ground beetle; Garden web worm; Grasshopper; Lygus bug; Pea weevil; Salt marsh caterpillar; Serpentine leaf miner; Bean leafhopper; Bean leaf roller; Corn earworm; <i>Compea surculio</i> .	2 qts. 1.1-3 pts.	Do not apply to green or snap beans after pods begin to form. No limitations for shelled beans for human consumption.
PEAS COWPEAS	Cutworm; Lygus bug; cowpea curculio; Darkling ground beetle; Southern green stink bug; Bean leafhopper; Bean leaf roller.	2 qts. 1.1-3 pts.	Do not apply after pods begin to form if pods are to be eaten for food or used for snags.
SOYBEANS	Armyworm; Bean leaf and Blister beetles; Corn earworm; Grasshopper; Green cloverworm; Velvet bean caterpillar; Cricket; Flea beetle; Thrips.	3 pts. 1 qt.	Do not apply less than 3 weeks before harvest.
BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS KOHLRABI	Flea beetle; Armyworm; Imported cabbageworm; Serpentine leaf miner; Thrips; Green peach aphid; Diamondback moth; Cabbage looper; Corn earworm; Cabbage aphid; Cutworm (climbing and surface feeding); Fall armyworm.	2 qts. 1.1-3 pts. 1 qt.	On broccoli, Brussels sprouts, cauliflower and shlrabi, do not apply after edible parts begin to form or within 30 days of harvest. On cabbage, do not apply after heads begin to form. On collards do not apply within 35 days of harvest.
CARROTS	Cutworms; Grasshopper; Flea beetle.	1.1-3 qts. 2 qts.	No time limitation. Apply as foliar spray only.
CELERY	Leafy leaf tier; Vegetable weevil; Cabbage aphid; Cutworms (climbing and surface feeding); Fall armyworms; Green apple aphid; Serpentine leaf miner; Thrips.	2.3-4 qts. 1 qt.	Do not apply after plants start to bunch or after plants are half mature.
CORN	Imported cabbageworm; Lesser cornstalk borer; Corn lantern fly; Armyworm; Chinch bug; Corn earworm; Grasshopper; Webworm.	1.1-3 qts. 2.3-4 qts. 1.1-3 qts.	No limitation on use of grain.
FRUIT	Corn billbug; Flea beetle.	1.1-3 qts.	Do not apply within 5 days of harvest.
SMALL GRAIN Barley, Oats, Rice Rye, Wheat	Armyworm; Beetles (blister, Colorado potato, cucumber, flea); cutworms (climbing and surface feeding); leafhopper; Serpentine leaf miner; Armyworm; Cutworm (climbing and surface feeding); Grasshopper; Chinch bug; Rice stink bug.	1.1-3 qts.	No restrictions on use of grain.
GRAIN SORGHUM	Chinch bug; Corn lantern fly; Cutworm; Grasshopper; Lesser cornstalk borer; Armyworm; Webworm; Corn billbug; Flea beetle.	2 qts. 1.1-3 qts. 1.1-3 qts.	Do not apply more than once after heads have begun to form. Do not apply within 28 days of harvest when using up to 1.1-3 qts. per acre or within 40 days when using 2 qts. per acre. Do not pasture dairy animals or animals being fattened for slaughter in treated fields. Do not ensile treated forage.

CROP
HORSERADISH
KALE
LETTUCE
OKRA
ONIONS
PARSNIPS
PEPPERS
PIMENTO
RUTABAGAS
STRAWBERRIES
TOMATOES

WARNING: Keep out of reach of children. Wash hands thoroughly after handling. Wash clothing. Wash if harmful if taken in physician. If spilled on skin, wash with water. Do not get in eyes. Do not get in mouth. Do not get in ears. DO NOT USE IN ENVIRONMENTAL CAUTION. This product is toxic if applied where runoff is likely. CONTAINER DISPOSAL: In a safe place. NON-WARRANTY: Foliar chemicals not specific. Buyer assumes responsibility.

SM 1 95

GENERAL INFORMATION:

Hi Yield TOXAPHENE 6 E.C. is an emulsifiable concentrate that must be diluted with water according to the directions below. It is intended for control of insect pests on agricultural crops. In these directions given below, Hi Yield TOXAPHENE 6 E.C. should be mixed with sufficient water to assure thorough coverage. Best control is obtained when application is made at first sign of insects or their damage appear. Applications should be repeated as necessary to maintain control.

GUIDE FOR APPROXIMATE AMOUNT OF WATER TO USE PER ACRE: For airplane and hydraulic sprayers 2 to 5 gallons; low gallonage sprayers 5 to 10 gallons. Use contrary to these directions may cause serious plant, crop or personal injury, excessive residues, unsatisfactory control or other unintended consequences.

NOTE: Do not feed treated vines, forage, plants, mill trash or ensilage made from plants to poultry, livestock, dairy animals or animals being finished for slaughter.

CROP	INSECTS	RATE PER ACRE	REMARKS
BEANS Green Snap lima	Armyworms; Bean beetles (Mexican bean, bean leaf, cucumber, wax bean, blister and flea); Bean leaf skeletonizer; Darkling ground beetle; Garden web worm; Grasshopper; Lygus bug; Pea weevil; Salt marsh caterpillar; Serpentine leaf miner;	2 qts.	Do not apply to green or snap beans after pods begin to form. No limitations for shelled beans for human consumption.
	Bean leafhopper; Bean leaf roller; Corn earworm; Cowpea curculio.	1 1/3 qts.	
PEAS COWPEAS	Cutworm; Lygus bug; cowpea curculio; Darkling ground beetle; Southern green stink bug;	2 qts.	Do not apply after pods begin to form if pods are to be used for snags.
	Bean leafhopper; Bean leaf roller.	1 1/3 qts.	
SOYBEANS	Armyworm; Bean leaf and Blister beetles; Corn earworm; Grasshopper; Green cloverworm; Velvet bean caterpillar; Cricket;	3 qts.	Do not apply less than 3 weeks before harvest.
BROCCOLI BRUSSELS SPROUTS CABBAGE CAULIFLOWER COLLARDS KOHLRABI	Flea beetle	1 qt.	On broccoli, Brussels sprouts, cauliflower and kohlrabi, do not apply after edible parts begin to form or within 30 days of harvest. On cabbage, do not apply after heads begin to form. On collards do not apply within 35 days of harvest.
	Armyworm; Imported cabbageworm; Serpentine leaf miner; Thrips; Green peach aphid; Diamondback moth; Cabbage looper; Corn earworm.	1 1/3 qts.	
CABBAGE COLLARDS KOHLRABI	Cabbage aphid; Cutworm (climbing and surface feeding); Fall armyworm.	1 qt.	No time limitation. Apply as foliar spray only.
	Cutworms; Grasshopper;	1 1/3 qts.	
CARROTS	Flea beetle	2 qts.	Do not apply after plants start to bunch or after plants are half mature.
CELERY	Celery leaf tier; Vegetable weevil;	2 3/4 qts.	
CORN	Cabbage aphid; Cutworms (climbing and surface feeding); Fall armyworms; Green apple aphid; Serpentine leaf miner; Thrips;	1 qt.	No limitation on use of grain.
	Imported cabbageworm;	1 1/3 qts.	
	Lesser cornstalk borer; Corn lantern fly;	2 3/4 qts.	
	Armyworm; Chinch bug; Corn earworm; Grasshopper; Webworm;	1 2/3 qts.	
EGGPLANT	Corn billbug; Flea beetle;	1 1/3 qts.	Do not apply within 5 days of harvest.
SMALL GRAIN Barley, Oats, Rice Rye, Wheat	Armyworm; Beetles (blister Colorado potato, cucumber leaf); cutworms (climbing and surface feeding); leafhopper; Serpentine leaf miner;	2 qts.	No restrictions on use of grain.
	Armyworm; Cutworm (climbing and surface feeding); Grasshopper; Chinch bug; Rice stink bug;	1 1/3 qts.	
GRAIN SORGHUM	Chinch bug; Corn Lantern fly; Cutworm; Grasshopper; Lesser cornstalk borer;	2 qts.	Do not apply more than once after heads have begun to form. Do not apply within 28 days of harvest when using up to 1 1/3 qts. per acre or within 40 days when using 2 qts. per acre. Do not pasture dairy animals or animals being fattened for slaughter in treated fields. Do not ensile treated forage.
	Armyworm; Webworm;	1 2/3 qts.	
	Corn billbug; Flea beetle;	1 1/3 qts.	

59.5 %
36.3 %
4.2 %
100.0 %

Crops and Small Grains
per Gallon

CHILDREN
Safety Warnings

GALLONS

CROP	INSECTS	RATE PER ACRE	REMARKS
PARSNIPS	Grasshopper;	2 qts.	Apply as a foliage treatment only. Do not use treated tops as food or feed.
KALE	Striped flea beetle;	1 qt.	
	Cabbage looper; Corn earworm; Diamondback moth;	2 2/3 qts.	Do not apply within 28 days of harvest when using 1 1/2 qts. per acre or within 35 days when using 2 2/3 qts. per acre.
Flea beetle;	2 qts.		
LETTUCE	Armyworm; Green peach aphid; Imported cabbage worm; Serpentine leaf miner; Thrips;	1 1/3 qts.	Do not apply after seeding stage on leaf lettuce or head lettuce after heads begin to form.
	Cutworm; Fall armyworm;	1 qt.	
	Cabbage looper; Cutworm; Field Cricket; Grasshopper; Salt marsh caterpillar; Flea beetle;	2 2/3 qts.	
	Armyworm; cucumber beetle;	1 1/3 qts.	
OKRA	Imported cabbageworm;	1 qt.	Do not use after pods begin to form.
ONIONS	Serpentine leaf miner;	2 qts.	
PARSNIPS	Thrips;	1 1/3 qts.	Do not apply to green or spring onions.
PEPPERS	Grasshopper; Striped flea beetle;	1 qt.	Apply to foliage only. Do not use tops for food or feed.
	Corn earworm; Cutworm; Fall armyworms; Flea hopper; Potato leafhopper; Red necked peanut worm; Southern green stink bug;	2 2/3 qts.	
PEPPERS	Velvetbean caterpillar;	2 qts.	Do not feed treated forage to dairy animals or animals being finished for slaughter.
	Thrips;	1 1/3 qts.	
PEPPERS	Armyworm; Beetles (blister Colorado potato, flea);	2 qts.	Do not apply within 5 days of harvest.
	Cutworm; Fruitworm; Hornworm; Pepper weevil; Serpentine leaf miner;	2 qts.	
PIMENTO	Flea beetle; Hornworm; Leaf miners; Pepper weevil;	1 1/3 qts.	For 1 1/3 qt. per acre rate, do not apply within 2 days of harvest.
RUTABAGAS	Flea beetle; Imported cabbageworm; Spinach leaf miner; Sugarbeet webworm;	2 2/3 qts.	Apply as a foliar treatment only. Do not use treated tops for food or feed.
SWEET POTATOES	Cutworm;	1 1/3 qts.	Do not apply after fruit starts to form.
	Cutworm; Strawberry crown borer; Strawberry weevil;	2 qts.	
TOMATOES	Armyworm; Blister beetle; Colorado potato beetle;	3 1/3 qts.	Do not apply within 3 days of harvest.
	Green chrysanthemum aphid; Hornworm; Pinworm; Russet mite; Serpentine leaf miner; Tomato fruit worm;	2 qts.	

WARNING: Keep out of reach of children. May be fatal if swallowed. Do not breathe spray mist. Do not get in eyes, on skin or on clothing. Wash thoroughly after use.

HARMFUL IF TAKEN INTERNALLY. Antidote: tablespoon of salt in a glass of warm water. Repeat until vomit fluid is clear. Call a physician.

If spilled on skin, wash thoroughly with soap and water. Wear respirator and goggles to protect from breathing spray mist or getting in eyes. Do not store near feed or foodstuffs. Do not feed treated forage to dairy animals or animals being finished for slaughter. Do not graze dairy animals or animals being finished for slaughter in field treated late in season. Do not apply after bolts open. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

ENVIRONMENTAL CAUTIONS: This product is toxic to bees and should not be applied when bees are actively visiting the area. This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from area treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

CONTAINER DISPOSAL: Do not reuse empty drum. Return to drum conditioner or destroy by perforating or crushing and burying in a safe place.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS

5M 1 95

EPA Reg. No. 7401-281 AA