

ACCEPTED

DEC 14 1972

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENT CIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 7401-65 SUBJECT TO ATTACHED COMMENTS.

1. WHAT IS THIS PACKAGE FOR?

We recommend this highly effective new fungicidal formulation, because it will make your lawn and garden disease control program easier.

2. HOW DO YOU USE IT?

SHAKE WELL BEFORE USE:

ROSES — For the control of Black Spot, Blossom Blight, mix 6 tablespoons in 1 gallon of water.

CHRYSANTHEMUMS — For the control of *Ascochyta* Ray Blight and Gray Mold.

GERANIUMS — For the control of Botrytis Blight (Leaf Spot):

IRIS — For the control of Botrytis Blight (Blossom Blight) and Iris Leaf Spot.

ZINNIA — For the control of Powdery Mildew:

*Mix 9 tablespoons in 1 gallon of water.

Spray plants on both sides of foliage to the drip point. Penetrate dense foliage. Apply at first sign of disease and repeat at 7-14 day intervals throughout growing season. Make application on a 7 day schedule during period of wet weather or severe disease conditions.

LAWNS — For the control of Brown Patch, Dollar Spot, Leaf Spot (Blight Going-out, Fading-out, and Melting-out), Gray Leaf Spot, Stem Rust on Bluegrass and Copper Spot mix 1 pint in 5-15 gallons of water and apply to 500 sq. ft. of lawn. The amount of water should be sufficient to provide complete coverage. DO NOT mow or water grass after treatment until the grass is thoroughly dry and preferably not for 24 hours after treatment. Start spray application in spring before diseases appear and repeat at 7-14 day intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Do not graze treated areas. Do not feed clippings to livestock.

VEGETABLES AND MELONS — Unless otherwise specified mix 3 tablespoons in 5-15 gallons of water and apply to 500 sq. ft. of garden area. Repeat at 7-10 day intervals or as necessary to maintain control. Carefully follow application schedule under individual crops.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER — For Alternaria Leaf Spot (S.E. States Only) and Downy Mildew. Apply after transplant, emergence of seeded crop or when conditions favor disease development.

CABBAGE — For Early and Late Blight. Begin when disease threatens.

POTATOES — For Botrytis Vine Rot, Early and Late Blight. Begin when plants are 6 inches high or disease threatens.

PUMPKIN, SQUASH (Summer and Winter), WATERMELON — For Anthracnose and Downy Mildew. Use 6 tablespoons per 5-15 gallons

of water per 500 sq. ft. for *Cercospora* Leaf Spot, Gum Stem Blight, Leaf Blight and Scab. Begin treatment in the first leaf stage.

TOMATOES — For Anthracnose, Early and Late Blights, Gray Leaf Spot, Gray Leaf Mold. Use 4 tablespoons in 5-15 gallons of water per 500 sq. ft. begin when disease threatens.

MELONS (Cantaloupe, Honeydew, Muskmelon) — For Anthracnose, Downy Mildew. Use 6 tablespoons in 5-15 gallons of water per 500 sq. ft. for Powdery Mildew (except for S.W. States), *Cercospora* Leaf Spot, Gummy Stem Blight, Leaf Blight, Scab. Begin when plants are in first true leaf stage or conditions favor disease development.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

ferti-lome BROAD SPECTRUM LIQUID FUNGICIDE will control listed fungi on roses, ornamentals, turf and lawn grasses, vegetables and melons thus permitting normal, vigorous growth for your enjoyment.

CAUTION: KEEP OUT OF REACH OF CHILDREN. May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Do not breathe spray mist. Wash hands, arms and face thoroughly with soap and warm water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse.

Do not contaminate feed or foodstuffs. Keep children and pets away from treated area until it is completely dry. Apply this product only as specified on this label.

This product is toxic to fish. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Food utensils such as tablespoons or measuring cups should not be used for food purposes after use with pesticides.

Do not reuse container for any purpose. Destroy empty container.

Product freezing temperature 32° F.**

**Satisfactory utility is restored after thawing and mixing of product. Repeated 0-10° freezing-thawing may impair utility.

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

*Daconil 2787 U.S. Patent No. 3,290,353 and No. 3,331,735. Diamond Shamrock Corp.

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS

28M-4-112

EPA Reg. No. 7401-65



ferti·lome



Broad Spectrum

LIQUID FUNGICIDE

- UNIQUE NEW FUNGICIDE FOR ROSES, ORNAMENTALS, TURF, LAWN GRASSES AND LISTED VEGETABLES AND MELONS.
- CONTROLS BLACK SPOT, BROWN PATCH, DOLLAR SPOT, LEAF SPOT, GRAY LEAF SPOT, BLOSSOM BLIGHT AND CERTAIN OTHER FUNGI
- 6 TABLESPOONS MAKE 1 GALLON OF SPRAY FOR ROSES.

CAUTION KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACCEPTED

DEC 14 1972

FOR PUBLICATION IN THE
 JOURNAL OF THE
 AMERICAN SOCIETY OF
 PLANT PATHOLOGISTS

1. WHAT IS THE PACKAGE FOR?
 A white plastic jug with a white cap and a white label. The label has a picture of a plant and some text.

2. HOW DO YOU USE IT?
SHAKE WELL BEFORE USE

ROSES — For Anthracnose, Downy Mildew, Gray Leaf Spot, Leaf Spot, and Stem Blight. Begin when plants are in the first leaf stage. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when disease first appears.

CHRYSANTHEMUMS — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Begin when plants are in the first leaf stage. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when disease first appears.

GERANIUMS — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Begin when plants are in the first leaf stage. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when disease first appears.

IRIS — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Begin when plants are in the first leaf stage. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when disease first appears.

ZINNIA — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Begin when plants are in the first leaf stage. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when disease first appears.

LAWNS — For Brown Patch, Dollar Spot, Leaf Spot, and Stem Blight. Apply when the grass is growing. Use 1/2 gallon of water per 100 sq. ft. Apply when the grass is growing.

VEGETABLES AND MELONS — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

CARROTS — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

POTATOES — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

PUMPKIN, SQUASH — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

WATERMELON — For Anthracnose, Downy Mildew, Gray Leaf Spot, and Stem Blight. Use 4 to 6 teaspoons in 1 1/2 gallons of water per 100 sq. ft. Begin when plants are in the first leaf stage.

VOLUNTARY PURCHASING GROUPS, INC.
 BONHAM, TEXAS

U.S. Pat. No. 3,440,000

EPA Reg. No. 2461-65



ferti·lome



'298

Broad Spectrum

LIQUID FUNGICIDE

- UNIQUE NEW FUNGICIDE FOR ROSES, ORNAMENTALS, TURF, LAWN GRASSES AND LISTED VEGETABLES AND MELONS.
- CONTROLS BLACK SPOT, BROWN PATCH, DOLLAR SPOT, LEAF SPOT, GRAY LEAF SPOT, BLOSSOM BLIGHT AND CERTAIN OTHER FUNGI.
- 6 TABLESPOONS MAKE 1 GALLON OF SPRAY FOR ROSES.

Active ingredient
 Chlorothalonil (tetrachloroisophthalonitrile)* 12.5%

Inert ingredients 87.5%

NET CONTENTS 16 FL. OZS. (ONE PINT)

CAUTION KEEP OUT OF REACH OF CHILDREN SEE BACK PANEL FOR ADDITIONAL CAUTIONS