(Front Panel)

1 A ? Pr. 51 Fr. 5 F 1 - 1 - 05

ACCEPTED with COMMENTS to EPA Letter Dated:

JUL2 5 1985

ferti-lome(R)

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Rag. No. 7401-65

٦

Broad Spectrum

Liquid Fungicide

^OUnique New Fungicide For Roses, Ornamentals, Turf, Lawn Grasses and Listed Vegetables and Melons.

(

;

^OControls Black Spot, Brown Patch, Dollar Spot, Leaf Spot, Gray Leaf Spot, Blossom Blight and Certain Other Fungi.

⁰6 Tablespoons Makes 1 Gallon of Spray For Roses.

| Active Ingredient: Chlorothalonil (tetrachloroisophthalonitrile)* | 12.5% |
|--|-------|
| Chlorothalonil (lettachlorothophilia) | 87.5% |
| Inert Ingredients: | 07.2% |

Keep Out Of Reach Of Children CAUTION See back panel for additional precautionary statements DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR? We recommended this highly effective new fungicidal - formula, because it will make your lawn and garden disease control program easier.

2. HOW DO YOU USE IT? SHAKE WELL BEFORE USE: ROSES - For control of Black Spot, Blossom Blight, mix 6 tablespoons in 1 gallon of water. CHRYSANTHEMUMS - For the control of Ascocyta Ray Blight and Gray Mold: GERANIUMS - For the control of Botrytis Blight (Leaf Spat). IRIS - For the control of Botrytis Blight (Plossom Blight) and Iris Leaf Spot: ZINNIA - For the control of Powdery Mildew: Mix 9 tablespoons in 1 gallon of water. Spray plants on both sides of foliage to the drip point. Penetrate dense folinge. Apply at first sign of disease and repeat at 7 - 14 day intervals throughout the growing season. Make application on a 7 day schedule during perioe of wet weather or severe disease conditions.

LAWNS - For the control of Brown Patch, Dollar Spot, Leaf Spot (Blight Going-out, Fading-out, and Melting-out), Gray Leaf Spot, Stem Rust on Bluegrass and Copper Spot mix 1 pint in 5 - 15 gallons of water and apply to 500 sq.ft. of lawn. The amount of water should be sufficient to provide complete coverage. DO NOT mow or water grass after treatment until the grass is thoroughly dry and preferable not for 24 hours after treatment. Start spray application in spring before diseases appear and repeat at 7 - 14 day intervals throughout the season. Make application on a 7 day schedule during periods of wet weather or severe disease conditions. Do not graze treated areas Do not feed clippings to livestock.

VEGETABLES AND MELONS - Unless otherwise specified mix 3 tablespoons in 5 - 15 gallons of water and apply to 500 sq.ft. of garden area. Repeat at 7 - 10 day intervals or as necessary to maintain control. Carefully follow application schedule under individual rops.

BROCCOLI, BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER - For Alternaria Leaf Spot (S.E. States Only) and Downy Mildew. Apply after transplant, emergence of seeded crop or when conditions favor disease development.

CARROTS - For Early and Late Blight. Begin when disease threatens. POTATOES - For Botrytis Vine Rot, Early and Late Blight. Begin when plants are 6 inches high or disease threatens.

PUMPKIN, SQUASH (Summer and Winter) WATERMELON - For Anthracnose and Downy Mildew. Use 6 tablespoons per 5 - 15 gallons of water per 500 sq.ft. for Cercospora Leaf Spot, Gum Stem Blight, Leaf Blight and Scab. Begin treatment in the first leaf stage. TOMATOES - For Anthracnose, Early and Late Blight, Gray Leaf Spot, Gray Leaf Mold. Use 4 tablespoons in 5 - 15 gallons of water per 500 sq.ft. Begin when disease threatens. MELONS (cantaloupe, Honeydew, Mushmelon) - For Anthracnose, Downy Mildew. Use 6 tables spoons in 5 - 15 gallons of water per 500 sq.ft. for Powdery Mildew (except For S.W. States), Cercospora Leaf Spot, Gummy Stem Blight, Leaf Blight, Scab. Begin when plants are in first true leaf stage or conditions favor disease development.

3. HERE ARE THE RESULTS YOU MAY EXPECT! ferti-lome Broad Spectrum Liquid Fungicide will control listed fungi on roses, ornamentals, turf and lawn grasses, vegetables and melons thus permitting normal, vigorous growth for your enjoyment.

BEST AVAILABLE COPY

STORAGE: Store in original container and place in a locked storage area. CONTAINER DISPOSAL: Do not reuse container. Wrap and put in trash.

PESTICIDE DISPOSAL Pesticide wastes are toxic. Improper disposal of excess pesticide, pesticide spray or rinsate is a folation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

NOTE TO USER: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reaction should contact a physician. Wear long sleeve shirt, long pants, and gloves while mixinc, loading and applying this product.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing spray mist. <u>Skin Sensitizer</u>, wash thoroughly after handling.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed - Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and Induce vomiting by touching back of throat with finger. Never induce vomiting or give anything by mouth to an unconscious person.

- If Inhaled Remove victim to fresh air. Apply artificial respiration if indicated.
- f On Skin Remove contaminated clothing. Wash affected areas with soap and water.
- If In Eyes Flush eyes with plenty of water. Get Medical attention if irritation persists.

NOTE TO PHYSICIAN: Persons having an allergic reaction, respond to treatment with antihistamines or storoid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in ne ghboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas.

Buyer assumes all risks of use, storage or handling of this materia: not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Product freezing temperature $32^{\circ}F$. Satisfactory utility is restored after thawing and mixing of product. Repeated 0 - $10^{\circ}F$ freezing - thawing may impair utility.

*Chlorothalon11 U.S. Patent 3,290,353 and No. 3,331,735

Voluntary Purchasing Groups, Inc.

Benham, Texas 75418

EPA Reg. No. 7401-65

У

. .

4.K.*

EPA Est. No. 7401-XT-1

5

Net Contents 32 fl. oz.