

Regno: 7401-66 PM-15. 191

1. WHAT IS THIS PACKAGE FOR?

This remarkable new insecticide controls over insects fast and for extended periods of time. It will greatly aid you with your Home Garden insect problems.

2. HOW DO YOU USE IT?

Mix 4 tablespoons in 3 gallons of water unless otherwise specified. Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage.

ROSES AND FLOWERS:

ROSES, AZALEAS, CAMELLIAS, RHODODENDRON, CHRYSANTHEMUMS, CARNATIONS, AND LILAC To kill Aphids, Cutworms, European Scale Crawlers, Flea Beetles, Lice Bugs, Leaf Miners (Dipterous), Leafhoppers, Mealybugs, Mites including Two-spotted Spider, European Red, Cyclamen, Carnation Bud and Shoot, Thrip^s and Whiteflies.

SHRUBS AND TREES:

AZALEA, CAMELLIA, HOLLY, ARBORVITAE, BIRCH, BOXWOOD, JUNIPER, RHODODENDRON, DOUGLAS FIR, ELM, LILAC, MIMOSA, OAK, PINE, POPLAR, SPRUCE and WILLOW. To kill Aphids, Mites (including European Red, Two-spotted Spider, Clover, Cyclamen and Privet), Webworms (Juniper and Mimosa), Scale Crawlers including Cottony-Cushion, Le canum, Pine Needle, Oak Pit Scale, San Jose and Soft, Whitemeles, Leaf Miners (including Arborvitae, Azalea, Birch, Boxwood, Dipterous, Holly and Oak), Leafhoppers, Mealybugs, Obscure Root Weevils and Omnivorous Leaf Tiers. For Mimosa Webworms apply a thorough foliar spray when adults first appear. Make two additional applications at 4-5 week intervals.


NOTE: Do not use on ferns, poinsettias, hibiscus, plumeras, gardenias and orchids since plant injury may occur.

LAWNS:

ST. AUGUSTINE, BERMUDA, BLUEGRASS, DICHONDRA, BENT, FESCUS, AND MARION To kill Sod Webworms, Chinch Bugs, Cutworms, Armyworms, Lawn Moths, Ants, Chiggers, Fleas, Earwigs, Pillbugs, Crickets, Mites (both Bermuda and Clover), Springtails, Ticks, Bermuda Grass Thrips, Frit Flies, Lawn Billbugs and Leafhoppers, water grass thoroughly a few hours before applying. Mix 4 tablespoons in 3 gallons of water and apply to 125 sq ft or 2 cups in 24 gallons of water to 1000 sq ft. NOTE: For lawn Billbugs, water lawn thoroughly immediately after application. In the north, Chinch Bugs will be killed using 2 tablespoons in 3 gallons of water per 125 sq ft.

FRUIT, CITRUS AND PECANS:

Mix 4 tablespoons in 3 gallons of water. Apply as a full cover spray unless otherwise specified. Assure thorough coverage. Repeat as necessary to maintain control.



ACCEPTED

MAY 1, 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 7401-66

DIAZINON®
LAWN & GARDEN

CROP	INSECTS	APPLY UP TO DAYS OF HARVEST	REMARKS
Peaches, Nectarines, Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Peach Twig Borer, Scale Crawlers, San Jose Scale Crawlers, Leafhoppers (peaches only), Pecan Mothlybugs.	20 10 10 = =	Apply as full cover spray. Apply when aphids, mites and crawlers first appear. For berries apply to peaches, in May and June. For leafhoppers (peaches only) apply as insectants develop. For "Pecan Mothlybug" apply as a cover spray from petals fall to 10%.
Apples	Leafhoppers, Eye-Spotted Bud Moths, Fruit Tree Leaf Rollers, Mealybugs	14 =	Apply as full cover spray. For Leafhoppers apply as insectants develop. For Eye-Spotted-Bud Moth, Fruit Tree Leaf Rollers and Mealybugs make 3-8 cover spray applications beginning in pink stage. Slight rusting may occur on some variety of apples, such as Golden Delicious.
Pear	Pear Psylla, Tanidorm Leaf Miners, Mealybugs, San Jose Scale Crawlers, Forbs Scale Crawlers	14	Apply as full cover spray for Pear Psylla mix 7 tablespoons in 3 gallons of water, spray at pre-pink stage. Make 1-2 additional applications during the first and second cover spray if necessary. For Tanidorm Leaf Miners and Mealybugs apply in 3 cover sprays, beginning in April.
Grapefruit Lemons Limes Oranges Tongelos Tangerines, Citrus Citron Kumquats and hybrids of these	Citrus Aphids, Fruit Tree Leaf Rollers, Cottony-Cushion Scale Crawlers, Soft Scale Crawlers, Citrus Thrips, Citrus Scale	21	For citrus aphids, Cottony-Cushion Scale Crawlers and soft scale crawlers and thrips, apply thorough cover spray when insects first appear. For Leaf Rollers apply when 50% of eggs in egg masses have hatched in spring. For citrus scale, apply as a postbloom spray and again as a summer-spray. For heavy soft scale infestations apply 8 tablespoons in 3 gallons of water. Thorough coverage of tree trunk, limbs and twigs is essential. Do not apply during bloom.
Pecans	Aphids, Mites, Pecan Nut Casebearers, Walnut Caterpillars		Apply as a thorough cover spray. Do not apply after husks crack.

VEGETABLES:

Mix 4 tablespoons in 3 gallons of water unless otherwise specified, apply to 1000 sq ft. Assure thorough coverage of upper and lower leaf surfaces. Repeat as necessary to maintain control.

Continued from page 2

CROP	INSECTS	REMARKS
Pole Beans	Black Bean Aphid, Mexican Bean Beetles, Leaf Miners	Do not apply within 7 days of harvest of beans. Beans and pea forage or bean and pea hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is cut for hay. Peas may be picked immediately.
Pea Beans	Dipterous Leaf Miners, Leafhoppers, Mites	
Peas	Aphids, Leaf Miners	
Cucumbers	Aphids, Serpentine Leaf Miners, Thrips	Do not apply within 7 days of harvest.
Tomatoes	Aphids, Dipterous Leaf Miners, Banded Cucumber Beetles, Fall and Southern Armyworms, Vinegar Flies (Drosophila spp.)	Do not apply within 1 day of harvest. For aphids and dipterous leaf miners mix 2 tablespoons in 3 gallons of water. For Aphid, Dipterous Leaf Miners and Vinegar Flies weekly applications are usually required.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

This new product, used as directed, will control listed insects in your lawn and garden.

WARNING: May be fatal if swallowed or absorbed through the skin. Do not breathe spray mist. Avoid contact