## 1. WHAT IS THIS PACKAGE FOR?

This product, developed with the reserior in mind, makes rose insect and disease control easier. It contains an insecticide-mitide and a fungicide that fills certain of your rose care needs.

### 2 HOW DO YOU USE 1T7

Mix 4 tablespoons in 1 gallon of water. Spray plants on both sides of follage to the drip point. Penetrate dense follage. Make new spray for each use. Follow directions as listed below.

ROSES: To kill Aphids, Mites (European Red and Two Spotted) Thrips, Leafhoppers and Dipterous Leaf Miners, to control Blackspot, Blossom Blight and to suppress Powdery Mildew, apply as pests of diseases appear, repeating application at 7-14 day intervals throughout growing season. Use 7 day intervals during wet weather or severe disease conditions.

CAMELIAS, AZALEAS, CHRYSANTHEMUMS, CARNATIONS, GLADIOLI, LILAC, RIODODENDRON, ARRONYITAE, BOX-WOOD, JUNIPER, HOLLY, EVERGREENS AND SMADE TREES: To kill Aphids, Mites, (including Clover, Cyclamen, European Red, Privet, Two-Spotted Spider, Carnotion Bud and Shoot), Bagworms, Leofhoppers, Leaf Miners (including Arborvitae, Azalea, Birch, Boxwood, Dipterous, Holly, and Ock), Whiteflies, Mealy Bugs, Scale Crawlers (including Cottony-Cushion, Lecanium, Pine Needle, Oak Pit Scale, San Jose, Soft and Euonymus Scale Crawlers), Thrips, Flea Beetles, Webworms (Juniper and Mimosa), European Pine Shoot Moths and Holly Bud Moths, apply as pests appear, repeating application as required. For Mimosa Webworms apply when adults agpear, repeating twice at 4-5 week intervals.

NOTE: Do not use on ferns, poinsettla, hibiscus, papaya, pilea, gardenia and orchids, since injury may occur.

# 3. HERE ARE THE RESULTS YOU MAY EXPECT!

When used as recommended ferti-lome ROSE SPRAY will kill the listed insects, control black spot and blossom blight, suppress powdery mildew on roses and kill the listed insects on other flowers and shrubs.

WARNING: May be fatal if swallowed or absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or household pets to go onto sprayed grass or come in contact with foliage until soray has compoletely dried.

# DO NOT USE, POUR, SPILL OR STORE NEAR HEAT O'R OPEN FLAME.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equpment, or disposal of waste.

This product is highly toxic to bees exposed to direct treatment of residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container for any purpose. Destroy empty container.

Food utensils such as tablespoons or measuring cups should not be used for food purposes after use with pesticides. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

\*Diazinon Reg. Trademark Geigy Agricultural Chemicals \*\*Daconil 2787 U. S. Pat. Nos. 3,290,353 and 3,331,735.

EPA Reg. No. 7401-67
VOLUNTARY PURCHASING GROUPS, INC.
BONHAM. TEXAS









# ACCEPTED

JAN 20 1972

UNDER THE FEDERAL INSECTICIDE
FUNCICIDE AND RODEWINCIDE ACT
FOR ECONOMIC POISON REGISTER.
TO ATTACHED COMMENTS.