

[FRONT PANEL]


ferti•lome®

# CARBARYL GARDEN SPRAY

[Optional proposed text appears in brackets - the final label may include some or all of the optional text]

For Homeowner Use Only

- [● Provides Control of Aphids, Thrips, Tent Caterpillars, Bagworms, Japanese Beetles, Fruit Tree Leaf Roller and Certain Other Insects on Roses, Ornamentals, Fruit and Citrus Trees]
- [● Kills Chinch Bugs, Armyworms, Ants and Tick in Lawns.]
- [● Controls San Jose Scale, Citricola Scale and Brown Scale Crawlers.]
- [● Contains Spreader Sticker.]



<b>ACTIVE INGREDIENT:</b>		
Carbaryl: 1-naphthyl N-methylcarbamate .....		25.0%
<b>OTHER INGREDIENTS:</b> .....		<u>75.0%</u>
<b>TOTAL:</b> .....		100.0%

**KEEP OUT OF REACH OF CHILDREN**

## CAUTION

**SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

**NET CONTENTS \_\_\_\_\_**

**[BACK/SIDE PANEL]**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply when weather conditions favor drift from area treated.

**1. WHAT IS THIS PACKAGE FOR?**

ferti•lome® Carbaryl Garden Spray was developed to provide you with an insecticide that will fill many of lawn and garden insecticide needs.

**2. HOW DO YOU USE IT?**

Mix 4 teaspoons in 1 gallon of water (for small jobs) or 8 oz. in 12 gallons of water or 1 pint in 24 gallons of water unless otherwise specified. Spray plants on both sides of foliage to the point of run off. Penetrate dense foliage. Make new spray for each use. Follow directions listed below.

**ROSES, AZALEAS, CAMELLIAS, GARDENIAS, CHRYSANTHEMUMS, RHODODENDRONS, ARBORVITAE, BOXWOOD, JUNIPERS, HOLLY, CARNATIONS, GLADIOLI, LILAC, EVERGREENS AND SHADE TREES:** To kill Rose, Elm Leaf and Apple Aphids; Rose Slugs, Birch Leaf, Boxweed and Oak Leaf Miners, Mealy Bugs, Leaf Rollers, Thorn, Plant, Lace and Boxwood Bugs, Bagworms, Elm Leaf, Blister, Japanese, June and Willow Leaf Beetles, Psyllids, Tent and Puss Caterpillars, Periodical Cicadas, Scale Insects, Knotweed, Orange Tortirix, Mimosa Webworms and Gypsy Moths, apply as pests appear, repeating application at 7-14 day intervals throughout growing season. NOTE: Do not apply to Boston Ivy since injury may occur.

**FRUIT AND CITRUS TREES:** Apply no less than 10 gallons of spray per mature tree as full coverage in normal spray schedule or when insects or their damage appear.

**PEACHES, APRICOTS, NECTARINES, PLUMS, PRUNES AND CHERRIES:** Mix 4 2/3 ozs. in 10 gallons of water for the control of Japanese Beetles, Lesser Peach Tree Borer, Peach Twig Borer, Plum Curculio, Fruit Tree Leaf Roller, Red-Banded Leaf Roller, Lecanium and San Jose Scale. For scale control apply when crawlers are present. For Lesser Peach Tree Borer control spray limbs and fruit thoroughly weekly during moth flight. Do not apply within 3 days of harvest.

**[BACK/SIDE PANEL CONTINUED]**

**GRAPEFRUIT, LEMONS, LIMES, ORANGES, TANGELOE, TANGERINES, CITRUS CITRUS CITRUS, KUMQUATS AND HYBRIDS OF THESE:** Use 6 2/3 ozs. in 10 gallons of water for the control of California Orangedog, Citrus Cutworms, Fruit Tree Leaf Roller, Orange Tortirix and Western Tussock Moth. Use 5 1/3 to 6 2/3 ozs. in 10 gallons of water for Black, Brown, Soft, California Red, Citrocola and Yellow Scale. Apply lower dosage early in season or against light infestation and the higher dosage late in season or against heavy infestations and situations involving infested fruit. Do not apply within 5 days of harvest.

**LAWNS:** To kill Chinch Bugs, Fall Armyworms, Leafhoppers, Sod Webworms or Lawn Moths, Ants, Earwigs, Fleas, Millepedes and Mosquitoes. Water lawn thoroughly. Mix 8 oz. (16 tablespoons) in 15 to 20 gallons of water and spray 675 sq. ft. of lawn. Repeat in 2-3 weeks if necessary. Do not allow children, pets or domestic animals on treated lawn until the insecticide is washed off the grass into the soil by sprinkling or rainfall and grass is completely dry.

For Brown Dog Ticks in lawns and dog's outdoor sleeping quarters, mix 1 pint in 3 gallons of water and spray inside and outside of doghouse and areas where ticks may hide when not feeding on dog. Place clean bedding in doghouse after spraying. Repeat in 2 weeks if necessary.

Not for direct application on dogs. Treat dog with an approved insecticide to prevent an infestation of lawn and sleeping quarters.

**3. HERE ARE THE RESULTS YOU MAY EXPECT!**

When used as recommended, this product will aid you in controlling damaging insects in your roses, flowers, ornamentals, fruit and citrus trees, and lawns.

**STORAGE AND DISPOSAL**

**STORAGE:** Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

**DISPOSAL: If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Harmful if swallowed or if absorbed through the skin. Avoid contact with skin, eyes, or clothing. Causes moderate eye irritation. Harmful if inhaled. Avoid breathing vapors or spray mist. Remove contaminated clothing and wash clothing before reuse. Wash thoroughly with soap and water after handling. Keep out of the reach of children and domestic animals.

**[BACK/SIDE PANEL CONTINUED]**

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAMES. Do not re-use container of any purpose. Destroy empty container. Do not contaminate feed or feedstuffs. Keep out of each of children. Keep children and pets away from treated area until area is completely dry. Apply this product only as specified on this label. Food utensils such as teaspoons or measuring cups should not be used for food purposes after use with pesticides.

**USER SAFETY REQUIREMENTS**

- \* Wear long-sleeved shirt, long pants, shoes plus sock and household latex or rubber gloves when mixing and applying this product.
- \* Wear a hat and eye protection when making overhead applications.
- \* Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
- \* Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- \* Do not allow children or pets to contact treated area until sprays have dried.

**FIRST AID****N-methyl Carbamate**

If swallowed	<ul style="list-style-type: none"> <li>• Call poison control center or doctor immediately for treatment advice.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not induce vomiting unless told to do so by the poison control center or doctor.</li> <li>• Do not give anything by mouth to an unconscious person.</li> </ul>
If inhaled	<ul style="list-style-type: none"> <li>• Move person to fresh air.</li> <li>• If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.</li> <li>• Call a poison control center or doctor for further treatment advice.</li> </ul>
If on skin or clothing	<ul style="list-style-type: none"> <li>• Take off contaminated clothing.</li> <li>• Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
If in eyes	<ul style="list-style-type: none"> <li>• Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li> <li>• Remove contact lenses if present, after the first 5 minutes, then continue rinsing eye.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-222-1222 or 1-800-900-4044 for emergency medical treatment information.

**NOTE TO PHYSICIAN:** Carbaryl is an N-methyl carbamate that inhibits cholinesterase. Atropine is an antidote. DO NOT use 2-PAM, opiates or cholinesterase-inhibiting drugs.

**ENVIRONMENTAL HAZARDS**

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or to areas where the surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of equipment washwater. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**NOTE:** BUYER ASSUMES ALL RISKS OF USE, STORAGE OR HANDLING OF THIS PRODUCT NOT IN STRICT ACCORDANCE WITH DIRECTIONS GIVEN HEREWITH. VOLUNTARY PURCHASING GROUPS, INC. WARRANTS THIS PRODUCT TO CONFORM TO THE CHEMICAL DESCRIPTION ON THE LABEL AND FOR THE PURPOSES STATED ON THE LABEL. THE LIMIT OF ANY LIABILITY INCURRED SHALL BE THE PURCHASE PRICE PAID BY THE USER OR BUYER.

Voluntary Purchasing Groups, Inc.  
P.O. Box 460  
Bonham, TX 75418

EPA Reg. No. 7401-83  
EPA Est. No. 7401-TX-1