

**1. WHAT IS THIS PACKAGE FOR?**

This new fungicide product developed to help you eliminate brown patch and other diseases in your lawns, flower beds and ornamentals.

**2. HOW DO YOU USE IT?**

**LAWNS** . . . nongrazed areas . . . brown patch (Rhizoctonia): Mix required dosage as listed below under your type of grass in recommended amount of water. Apply to specified area to be treated by means of a sprinkler can, hose-on sprayer or any other convenient method of obtaining even coverage. For all types of grass sprinkle lightly after application to insure movement of chemical through grass to the soil level. NOTE: Under certain growing conditions, a temporary discoloration of the grass occasionally may occur.

**ST. AUGUSTINE, BERMUDA AND OTHER SIMILAR TYPES OF WARM SEASON GRASSES:** Mix 6 tablespoons in 1 gallon of water to treat 100 sq. ft. For larger area mix 1 quart in 15 gallons of water for 1000 sq. ft. Spray in fall or spring repeating application in 3-4 weeks.

**BLUEGRASS, FESCUES AND OTHER COOL SEASON GRASSES SUCH AS RYEGRASS:** Mix 18 tablespoons (9 oz.) in 4 gallons of water to treat 1000 sq. ft. Begin treatment before disease is expected to appear, repeating at 7-10 day intervals during hot weather.

**DICHONDRA:** Mix 10 tablespoons (5 oz.) in 1 gallon of water and apply to 100 sq. ft. For larger area mix 1 1/2 quarts in 40 gallons of water and apply to 1000 sq. ft. Apply in fall or spring at first sign of disease and repeat 3-4 weeks later.

**AZALEAS** . . . petal blight (Ovulinia), **CAMELIAS** . . . flower or petal blight (Sclerotinia): Mix 32 tablespoons (16 oz.) in 10 gallons of water and apply to 50 sq. ft. Spray ground and surrounding area under plants. Begin 4 weeks before bloom. Repeat at 3-4 week intervals through bloom period.

**POINSETTIAS, CHRYSANTHEMUMS, BEGONIAS, AFRICAN VIOLETS, CARNATIONS, SNAPDRAGONS AND SIMILAR BEDDING PLANTS** . . . stem rot (Rhizoctonia): Mix 11 1/2 tablespoons (5 3/4 oz.) in 30 gallons of water and apply to 50 sq. ft. Use as a soil drench before planting. **CALENDULA, LARKSPUR, SNAPDRAGON AND SWEET PEA** . . . stem rot (Sclerotinia): Mix 15 tablespoons (7 1/2 oz.) in 25 gallons of water and apply to 50 sq. ft. Use as a soil drench 1 week before planting.

**3. HERE ARE THE RESULTS YOU MAY EXPECT!**

When used as directed this new product will rid your lawn of unsightly brown patch and control other fungal diseases in your flower and ornamentals.

**CAUTION:** Keep out of reach of children. May be harmful if swallowed. Do not take internally. Avoid breathing spray mist. Avoid contact with eyes and skin. May cause skin irritation. Wash thoroughly after using.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Food utensils such as tablespoons or measuring cups should not be used for food purposes after use with pesticides.

Do not graze treated areas. Do not feed clippings to livestock.

Do not contaminate feed and foodstuffs. Do not reuse container for any purpose. Destroy empty container.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where run-off is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of waste. Apply this product only as specified on this label.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**VOLUNTARY PURCHASING GROUPS, INC.**

Bonham, Texas

28M-3-114

EPA Reg. No. 7401-84-AA



ferti·lome



LIQUID FUNGICIDE FOR

**Brown Patch Control**

5498

**CAUTION** KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS

EPA Est. 7401-TX-1

ACCEPTED

APR 23 1975

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC POISON REGISTERED UNDER NO. 7401-84

**1. WHAT IS THIS PACKAGE FOR?**

This new fungicide product developed to help you eliminate brown patch and other diseases in lawns, flower beds and ornamentals.

**2. HOW DO YOU USE IT?**

**LAWNS** - Nongrazed areas - brown patch (Rhizoctonia) - Mix required dosage as listed below under your type of grass in recommended amount of water. Apply to soil. Let area to be treated by means of a sprinkler, can, hose or sprayer or any other convenient method of obtaining even coverage. For all types of grass, sprinkle lightly after application to insure movement of chemical through grass to the soil level. **NOTE:** Under winter growing conditions, soil temperature is a clue as to whether the grass is or is not dormant.

**ST. AUGUSTINE, BERMUDA AND OTHER SIMILAR TYPES OF WARM SEASON GRASSES** - Mix 6 tablespoons (2 oz.) of water to treat 100 sq. ft. For larger areas mix 1 quart (16 gallons of water for 1000 sq. ft.) Spray or filter spray on repeating application in 4 weeks.

**RYEGRASS, FESCUE AND OTHER COOL SEASON GRASSES SUCH AS FINE FESCUE** - Mix 12 tablespoons (2 cups) of water to treat 100 sq. ft. Begin treatment before disease is expected to appear, repeat at 2-4 week intervals. Best for warm weather.

**DORMANT LAWNS** - Mix 12 tablespoons (2 cups) of water to treat 100 sq. ft. For larger areas mix 1 quart (16 gallons of water) and apply to 1000 sq. ft. Apply in late autumn or early spring. Repeat in 4 weeks.

**AZALEAS, CAMELLIAS AND OTHER ORNAMENTALS** - Flower or petal blight (Sclerotinia) - Mix 2 tablespoons (1/2 oz.) in 10 gallons of water and apply to 50 sq. ft. spray ground and surrounding area under plants. Begin 4 weeks before bloom. Repeat at 2-4 week intervals through bloom period.

**POINSETTIAS, CHRYSANTHEMUMS, BEGONIAS, AFRICAN VIOLETS, CANNATIONS, SNAPDRAGONS AND SIMILAR BEDDING PLANTS** - Stem rot (Rhizoctonia) - Mix 1 1/2 tablespoons (1/2 oz.) in 30 gallons of water and apply to 50 sq. ft. Use as a soil drench before planting.

**PEONIES, LARKSPUR, SNAPDRAGON AND SWEET PEAS** - Stem rot (Sclerotinia) - Mix 1 1/2 tablespoons (1/2 oz.) in 30 gallons of water and apply to 50 sq. ft. Use as a soil drench 1 week before planting.

**3. HERE ARE THE RESULTS YOU MAY EXPECT:**

When used as directed this new product will rid your lawn of unsightly brown patch and control other fungal diseases in your flower and ornamentals.

**CAUTION:** Keep out of reach of children. May be harmful if swallowed. Do not take internally. Avoid breathing spray mist. Avoid contact with eyes and skin. May cause skin irritation. Wash thoroughly after using.

**DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.**

Food utensils, such as tablespoons or measuring cups should not be used for food purposes after use with product.

Do not spray treated areas. Do not feed plants to livestock.

Do not use on lawns, lawns, and feedstuffs. Do not reuse container for any purpose. Destroy empty container.

Do not use on lawns, lawns, and feedstuffs. Do not reuse container for any purpose. Destroy empty container. Do not use on lawns, lawns, and feedstuffs. Do not reuse container for any purpose. Destroy empty container.

Do not use on lawns, lawns, and feedstuffs. Do not reuse container for any purpose. Destroy empty container.

**VOLUNTARY PURCHASING GROUPS, INC.**

Bonham, Texas

EPA Reg. No. 74-1-14 AA



**LIQUID FUNGICIDE FOR** \$4<sup>98</sup>

# Brown Patch Control

- Controls brown patch (Rhizoctonia) in warm and cool season grasses such as St. Augustine, Bermuda, Bluegrass, Fescue, Ryegrass, Dichondra and similar type grasses
- Ends circular brown spots of dead grass in lawns
- Controls listed diseases on Azaleas, Camellias and other listed flowers and ornamentals
- Induces healthy lawn, flower and shrub growth thru disease control
- 6 Tablespoons make 1 gallon of spray
- 1 Quart treats 1000 sq. ft. of lawn

Active Ingredient: Pentachloronitrobenzene .....	24%
Inert Ingredients .....	76%
Total 100%	

**CAUTION KEEP OUT OF REACH OF CHILDREN**  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

NET CONTENTS 32 FL. OZS.  
(ONE QUART)

EPA Est. 7401-TX-1

**ACCEPTED**

UNDER THE FEDERAL INSECTICIDE  
FUNGICIDE AND RODENTICIDE ACT  
FOR ECONOMIC POISON REGISTERED  
UNDER NO.