

3/3

STORAGE: Store in original container and place in a locked storage area.

CONTAINER DISPOSAL: Do not reuse container. Wrap and put in trash.

PESTICIDE: Improper disposal of excess pesticide, pesticide spray or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

NOTE TO USER: This product may product temporary allergic side effects characterized areas. Persons having allergic reaction should contact a physician. wear long sleeve shirt, long pants, and gloves while mixing, loading and applying this product.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, aquatic invertebrates, and marine/estuarine organisms. Runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water. Do not apply when weather conditions favor drift from treated areas. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

*Diazinon Reg. Trademark of Ciba-Geigy Chemical CO.

** Daconil US Patent No. 3,290,353 and 3,331,735


Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-67

Net Contents 16 fl. oz.



7401-86 PM 15
19 1 of 3
THIS PRODUCT MUST NOT BE USED ON GOLF COURSES OR SOD FARMS
(Front Panel)

WORM SPRAY

ROSES, FLOWERS, ORNAMENTALS, LAWNS, TREES AND SHRUBS
AND VEGETABLES

* CONTAINS SPREADER-STICKER

* 4 TABLESPOONS MAKE 1 GALLON OF SOLUTION

BEST AVAILABLE COPY

ACTIVE INGREDIENTS:

0,0-Dimethyl (1,1-Dimethyl-2,2-bis(4-methylphenyl)ethane)

2,4-Dichlorophenoxyacetic acid

*INERT INGREDIENTS..... 87.50%

*(Contains 77.5% Aromatic Petroleum Derivatives)

Total . . . 100.00%

WARNING (18 Point)

KEEP OUT OF REACH OF CHILDREN (12 Point)

SIDE PANEL FOR ADDITIONAL WARNINGS (12 Point)

(Side Panel)

1. WHAT IS THIS PACKAGE FOR?

This package is the first specific worm spray introduced into the Home and Garden market. It was developed to make it easy to kill worms quickly and maintain worm control.

2. HOW DO YOU USE IT?

Mix 4 tablespoons in 3 gallons of water unless otherwise specified. Spray plants on both sides of foliage to the point of runoff. Penetrate all dense foliage. Repeat as necessary to maintain control. Use a spray for each use. Follow directions listed below.

ROSES, FLOWERS, ORNAMENTALS, TREES AND SHRUBS: To kill Cutworms, Bagworms, Juniper and Mimosa Webworms, apply as directed above. For Mimosa Webworms apply as thorough foliar spray when adults first appear. Make two additional applications at 4 - 5 week intervals.

For Tent Caterpillars and Caneborer and Fall Webworm: mix 4 tablespoons in 3 gallons of water. Apply as a thorough spray.

LAWNS: May be used on St. Augustine, Bermuda, Bent Bluegrass, Fescue, Merion and Dichondra. To kill Armyworms, Cutworms and Sod Webworms mix 4 tablespoons in 3 gallons of water and apply to 125 sq. ft. Thoroughly water grass a few hours before applying. Repeat as necessary to maintain control.

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

7401-86

(Side Panel)

283

In three gallons of water and cover approximately 1000 sq. ft. of garden. Thoroughly spray all plant parts. Apply when insects first appear. Repeat as necessary to maintain control. Do not spray on windy days.

TOMATOES. To kill Fall and Summer Armyworms. Do not apply 7 day of harvest.

CABBAGE, BROCCOLI, and CAULIFLOWER. To kill Imported Cabbage worm. Do not apply to broccoli or cauliflower within 5 days of harvest. Do not apply to cauliflower within 12 days of harvest.

SWISS CHARD. To kill Imported Cabbage worm. Do not apply to Swiss chard within 12 days of harvest.

POTATOES. To kill Southern Armyworms. Do not apply within 14 days of harvest.

SWEET CORN. To kill Corn Earworms. Mix 5 tablespoons (2 1/3 ozs.) in 3 gallons of water and apply to 1000 sq. ft. when silks first appear. Two or three repeat applications at 2 - 3 day intervals will be necessary. Corn may be picked immediately following last application.

3. HERE ARE THE RESULTS YOU MAY EXPECT!
This new worm spray will quickly kill destructive worms found in your shrubs, gardens, lawns and vegetables, and give long lasting control.

WARNING: May be fatal if swallowed or absorbed through the skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of feed and water. Do not permit children or household pets to go on sprayed grass or come in contact with foliage until spray has completely dried. Do not store or use near heat or open flame.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agriculture Extension Service.

BEST AVAILABLE COPY

