

WHAT IS IT? Spider mites, the almost invisible yet one of the most destructive insects found in your Home Garden, can be controlled with this new formulation. It offers you quick acting long lasting control of mites and other listed insect pests.

HOW DO YOU USE IT?

To kill Spider mites, Bagworms, Aphids and other listed insects, follow dilution rates under plant headings. Make adequate sprays when pests appear or when their damage is first observed. Do not over-dose or over-spray. Do not use on ornamental plants not listed. The uses listed below are foliar sprays unless otherwise indicated.

ROSES, AZALEA, CAMELLIAS, GARDENIAS, ARBORVITAE, HOLLY, BOXWOOD, HEMLOCK, POINSETTIAS, GLADIOLUS AND GERBERAS: Mix 2 tablespoons in 1 gallon of water. Apply as directed above.

ROSES: Mites, Aphids, Leafhoppers and Thrips.
AZALEA: Mites, Tea Scale, Leaf Miners, Lace Bugs and Whiteflies.

CAMELLIAS: Mites, Aphids, Camellia and Tea Scale.

Foliar Spray: Apply 2 sprays 6 weeks apart the first year followed by annual applications soon after first new growth begins in spring.

Soil Drench: Apply as a soil drench around base of plants in early spring at the rate of 12 tablespoons per 1 gallon of water per plant up to 6 feet tall. Increase the rate proportionately for larger plants.

GARDENIAS: Tea Scale and Whiteflies.

ARBORVITAE: Mites, Bagworms and Aphids.

HOLLY: (English and American) not Burford variety. Mites, Leaf Miners and Soft Scale. For Leaf Miners apply in spring when leaf miner flies first appear or in early summer for control of larva in infested areas.

BOXWOOD: Mites, Leaf Miners and Mealy Bugs. (For leaf miners apply as under HOLLY).

HEMLOCK: Mites and Scale.

POINSETTIAS: (Outdoor plants) Mites, Aphids, Mealy Bugs and Whiteflies.

GLADIOLUS: Aphids and Thrips.

GERBERAS: Thrips.

JUNIPER, EUONYMUS, PINES, YEWs, CEDAR, OAK, IRIS and DAY LILLIES: Mix 4 tablespoons in 1 gallon of water. Apply as directed above.

JUNIPER: Mites, Bagworms, Aphids and Midges.

EUONYMUS: Aphids and Scale.

PINES: Bagworms, Aphids, European and Nantucket Pine Tip Moth and Zimmerman Pine Moth.

YEWs: (Taxus) Mites, Mealy Bugs and Fletcher Scale.

CEDAR: Mites.

OAK: Golden Oak Scale.

IRIS: Aphids, Iris Borers and Thrips. For borer control, spray when new leaves are 5-6 inches tall.

DAY LILLIES: Aphids and Thrips. NOTE: Do not use on Easter Lilies as injury may result.

FRUIT TREES: To kill listed insects mix required dosage in following table in 1 gallon of water. Apply as directed below. Do not apply during bloom period. Do not use on citrus seedlings. Make no more than 2 applications to mature fruit.

| | |
|--------------------|---|
| APPLES AND PEARS | Mix 1-2 tablespoons per gallon for Aphids, Leafhoppers, Pear Psylla and Mites (except rust mites). Do not apply within 28 days of harvest. Apply as a thorough distribution coverage spray. |
| LEMONS AND ORANGES | Mix 1-2 tablespoons per gallon for Aphids. Do not apply within 15 days of harvest. Apply as an outside coverage spray. |
| | Mix 1-2 tablespoons per gallon for Mites (except rust mites). Do not apply within 15 days of harvest. Apply as a thorough distribution coverage spray. |
| | Mix 2-3 tablespoons per gallon for Scales (except black and snow). Do not apply within 45 days of harvest. Apply as a thorough coverage spray. |
| | Mix 1-2 tablespoons per gallon for Thrips. Do not apply within 15 days of harvest. Apply as a mist spray. |
| | Mix 2 tablespoons per gallon for White Flies. Do not apply within 15 days of harvest. Apply as a thorough distribution coverage spray. |

When used as directed ferti-lome Systemic SPIDER MITE SPRAY will kill and give long lasting control of spider mites that have invaded your ornamental garden areas.

CAUTION: KEEP OUT OF REACH OF CHILDREN. VAPOR HARMFUL IF SWALLOWED. CONCENTRATED MATERIAL CAUSES EYE IRRITATION. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not take internally. Avoid contact with skin, eyes and clothing. In case of contact with eyes flush with plenty of water for at least 15 minutes, get medical attention. Do not breathe vapors or spray mist. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling.

Food utensils such as a tablespoon or measuring cup should not be used for food purposes after use with pesticides.

TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse container for any purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon.

*CYGON Reg. Trademark of American Cyanamid Company.

14M-1-101

EPA Reg. No. 7401-106

VOLUNTARY PURCHASING GROUPS, INC., BONHAM, TEXAS



systemic

SPIDER MITE SPRAY

- KILLS DESTRUCTIVE SPIDER MITES: CAMELLIAS, ARBORVITAE AND OTHER SHRUBS, SHADE TREES AND FRUIT TREES.
- KILLS BAGWORMS, APHIDS, THRIPS, AND MANY OTHER ORNAMENTAL GARDEN PLANTS.
- NEW TRIPLE ACTION INSECTICIDE: CONTACT AND RESIDUAL INSECTICIDE ACTION.
- 2 TABLESPOONS MAKE 1 GALLON OF SPRAY.

NET CONTENTS ONE PINT
Active Ingredient: Dimethoate (O,O-dimethyl S-(N-methylcarbamoyl phosphorodithioate)*
Inert Ingredients: Total

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL PRECAUTIONS.

ACCEPTED WITH COMMENTS

ACCEPTED

| | |
|--------------------|--|
| APPLES AND PEARS | Mix 1-2 tablespoons per gallon for Aphids, Leafhoppers, Pear Psylla and Mites (except rust mites) Do not apply within 28 days of harvest. Apply as a thorough distribution coverage spray. |
| LEMONS AND ORANGES | Mix 1-2 tablespoons per gallon for Aphids. Do not apply within 15 days of harvest. Apply as an outside coverage spray. |
| | Mix 1-2 tablespoons per gallon for Mites (except rust mites) Do not apply within 15 days of harvest. Apply as a thorough distribution coverage spray. |
| | Mix 2-3 tablespoons per gallon for Scales (except black and snow) Do not apply within 45 days of harvest. Apply as a thorough coverage spray. |
| | Mix 1-2 tablespoons per gallon for Thrips. Do not apply within 15 days of harvest. Apply as a mist spray. |
| | Mix 2 tablespoons per gallon for White Flies. Do not apply within 15 days of harvest. Apply as a thorough distribution coverage spray. |

When used as directed, ferti-lome Systemic SPIDER MITE SPRAY will kill and give long lasting control of spider mites that have invaded your ornamental garden areas.

CAUTION: KEEP OUT OF REACH OF CHILDREN. VAPOR HARMFUL. HARMFUL IF SWALLOWED. CONCENTRATED MATERIAL CAUSES EYE IRRITATION. DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

Do not take internally. Avoid contact with skin, eyes, and clothing. In case of contact with eyes, flush with plenty of water for at least 15 minutes; get medical attention. Do not breathe vapors or spray mist. Avoid contamination of feed and foodstuffs. Wash thoroughly after handling.

Food utensils, such as a tablespoon or measuring cup, should not be used for food purposes after use with pest codes.

TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not contaminate water by cleaning of equipment or disposal of waste.

Do not reuse container for **any** purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given hereon with.

*CYGON Reg. Trademark of American Cyanamid Company.

14M 1-101 EPA Reg. No. 1401-101

VOLUNTARY PURCHASING GROUPS, INC., BONHAM, TEXAS



systemic

SPIDER MITE SPRAY



\$2.98

- KILLS DESTRUCTIVE SPIDER MITES ON ROSES, AZALEAS, CAMELLIAS, ARBORVITAE AND OTHER LISTED ORNAMENTAL SHRUBS, SHADE TREES AND FRUIT TREES
- KILLS BAGWORMS, APHIDS, THRIPS, SCALE INSECTS AND CERTAIN OTHER ORNAMENTAL GARDEN INSECTS ON CERTAIN PLANTS
- NEW TRIPLE ACTION INSECTICIDE PROVIDES SYSTEMATIC, CONTACT AND RESIDUAL INSECT CONTROL
- 2 TABLESPOONS MAKE 1 GALLON OF SPRAY

NET CONTENTS ONE PINT LIQUID

| | |
|---|------|
| Active Ingredient | |
| Dimethoate (O,O-dimethyl S-(N-methylcarbamoylmethyl) phosphorodithioate)* | 8% |
| Inert Ingredients | 92% |
| Total | 100% |

CAUTION: KEEP OUT OF REACH OF CHILDREN. SEE BACK PANEL FOR ADDITIONAL CAUTIONS.