

Not Reviewed. Registrant claims to be in accordance with Reregistration of Pesticide Product Guidance for Phase 2 Response, Page 2.12, Active to Inert Change in Status.

7401-115 1A10
1 10 1

(FRONT PANEL)

HI-YIELD

8 LB. MALATHION

EMULSIFIABLE CONCENTRATE

(Contains 8 lbs. of Malathion per gallon)

ACTIVE INGREDIENTS:

Malathion* 81.4%

**INERT INGREDIENTS: 18.6%

**(Contains 4.0 Aromatic Petroleum Derivatives)

Total 100.00%

* O,O-dimethyl dithiophosphate of diethyl mercaptosuccinate

CAUTION (18 POINT)

KEEP OUT OF REACH OF CHILDREN (12 POINT)

HARMFUL IF SWALLOWED. AVOID BREATHING SPRAY MIST. AVOID CONTACT WITH SKIN, EYES AND CLOTHING. WASH THOROUGHLY AFTER HANDLING. AVOID CONTAMINATION OF FEED AND FOODSTUFFS. THIS PRODUCT IS TOXIC TO FISH AND WILDLIFE. KEEP OUT OF ANY BODY OF WATER. KEEP CONTAINER CLOSED AND AWAY FROM HEAT, SPARKS AND OP FLAME.

DO NOT RE-USE CONTAINER, DESTROY WHEN EMPTY.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS

NET CONTENTS GALLONS

GENERAL DIRECTIONS

Hi-yield 8 lb. MALATHION EMULSIFIABLE CONCENTRATE (8lb. per gallon) may be used in conventional hydraulic sprayers and low volume applicators. Mix this product with sufficient water for uniform coverage. Repeat application as necessary. Use this product only for recommended purposes and at recommended dosages. Follow state and federal recommendations.

USE DIRECTIONS

CROP	INSECT	DOSAGE	DIRECTIONS
Cotton	Boll weevil	1-2 pts./acre	Use when insect is resistant to other insecticides. Apply on foliage every 3-7 days until controlled. No time limitation.
	Brown Cotton Leafworm	1/4 pt./acre	
	Cotton aphid	1/2-1pt./acre	
	Cotton flea-hopper	1/2-2 1/2 pts./acre	
	Cotton leaf perforator	1 pt./acre	

7401-115

CROP	INSECT	DOSE	DIRECTIONS
Kohlrabi	Aphids	1 1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Lettuce, Outdoor	Aphids	1-2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 14 days to harvest (if for seed). Allow 7 days to harvest if for leaf.
Pea (black-eyed)	Aphids	1/2 pts./acre	Apply on foliage as needed. Allow 7 days to harvest if forage is to be fed to livestock.
Pea (green)	Aphids	1/2 pts./acre	Apply to foliage during early bloom period as soon as eggs are visible. Allow 7 days to harvest.
Pea (wax)	Aphids	1/2 pts./acre	Apply on foliage as needed. Allow 7 days to harvest if forage is to be fed to livestock.
Pea (other)	Aphids	1/2 pts./acre	Apply on foliage as needed. Allow 7 days to harvest if forage is to be fed to livestock.
Turnip (field)	Drosophila	2 pts./acre	Apply on foliage as needed. Allow 1 day to harvest.
Turnip (field)	Spider mites	1 1/2 pts./acre	Apply on foliage as needed. Allow 1 day to harvest.
Turnip (field)	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.

BEST AVAILABLE COPY

CROP	INSECT	DOSAGE	DIRECTIONS
Rice	Stink bug	3/4-lpt./acre	Apply about 1 week after heads appear. Two applications may be needed.
Sugar Beets	Grasshoppers	1-2 pts./acre	Allow 7 days to harvest. Apply on foliage and soil as needed. Allow 7 days to harvest.
Beet, White	Foliage leathopper	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Beet, White	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Kale	Spider Mites	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.
Kale	Aphids	1/2 pts./acre	Apply on foliage when aphids appear; repeat weekly as needed. Allow 7 days to harvest.

BEST AVAILABLE COPY