

GENERAL INFORMATION: HI-YIELD DSMA LIQUID HERBICIDE is used for selective post-emergent weed control, particularly for grassy weeds. It is a combination of herbicide and surfactant. It is unnecessary to add any other surfactant to the spray solution. Best results are obtained on young, actively growing weeds at air temperatures above 70° F.

MIXING INSTRUCTIONS: HI-YIELD DSMA LIQUID HERBICIDE must be thoroughly dissolved. Fill the spray equipment reservoir one-half full with water and add the required amount of herbicide with agitation. Finish filling the reservoir with water and apply. After use, clean equipment thoroughly by flushing with water. Do not store spray solution in tank for a prolonged period.

CONTAINER DISPOSAL: Do not reuse empty container. Wash thoroughly with detergent, crush if possible, and discard in a safe place.

DIRECTIONS FOR USE:

COTTON: HI-YIELD DSMA LIQUID HERBICIDE is useful as a directed application in cotton for post-emergent control of weeds and grasses such as Johnsongrass, nutsedge, Dallisgrass, cocklebur, barnyardgrass, ragweed, sandbur and puncturevine.

Add 1 gallon of HI-YIELD DSMA LIQUID HERBICIDE to 40 gallons of water and mix thoroughly. Do not add any surfactant to the spray solution. Apply spray solution at 33 gallons per acre. Weed control is most effective at temperatures above 70° F. For band application in 40 inch rows, apply 1 gallon of spray solution per inch of band width per acre. Direct the spray solution to the base of the cotton plant. Nozzles should be placed so as to avoid spraying the cotton foliage. A second application may be required during rapid growing conditions to control perennials such as Johnsongrass.

Apply the second application 1 to 3 weeks later.

When treated, a reddish discoloration of the cotton plant will develop normally.

APPLY ONLY WHEN COTTON IS 3 INCHES HIGH TO FIRST BLOSSOM. DO NOT APPLY AFTER FIRST BLOSSOM. DO NOT FEED TREATED FOLIAGE TO LIVESTOCK IN GRAZE TREATED AREAS.

CITRUS (bearing): HI-YIELD DSMA LIQUID HERBICIDE is used for control of Johnsongrass, Dallisgrass, sandbur (sandspur), nutsedge, barnyardgrass, chickweed, wood sorrel, cocklebur, puncturevine and other similar weeds in citrus orchards (grapefruit, lemons, limes, oranges, tangerines). NOTE: Not for use in Florida.

11-1-72

7401-130