

DIRECTIONS: For Use Against Certain Insects Infesting Range Cattle and other stock.

PRECAUTIONS IN USING THIS PRODUCT: Do not treat animals under one month of age. When applying sprays, avoid contamination of feed, food containers and watering troughs. DO NOT APPLY TO DAIRY ANIMALS.

NOTE: Do not use this product as a dip treatment. Use only as a spray.

ANIMAL	PEST	RATE IN WATER	REMARKS
Beef Cattle (Range Cattle)	Lice	2 gal. per 100 gal. or 13 ozs. per 5 gal.	Treat animals thoroughly. Repeat application in 10 days and thereafter as needed.
	Ticks	2 2/3 gal per 100 gal. or 17 ozs. per 5 gal.	Treat animals thoroughly. Repeat application at 2 week intervals as needed.
	Hornflies	2/3 gal. per 100 gal. or 13-20 ozs. per 5 gal.	Apply on back and neck of animals and repeat appli- cation at 10-14 day inter- vals.
Hogs	Lice	2 gal. per 100 gal.	Apply a complete coverage spray to animals (includ- ing the inside of ears), pens and litter. Repeat in 10 days and thereafter as needed. Sprayed hogs should be kept out of sun and wind for a few hours after treatment.
	Sarcoptic Mange	2 2/3 gal per 100 gal. or 17 ozs. per 5 gal.	
Sheep & Goats	Lice Ticks Keds	2 2/3 gal per 100 gal. or 17 ozs. per 5 gal.	Spray animals thoroughly. Repeat application in 10 days and thereafter as needed.

Hi-Yield. M. & M. RANGE CATTLE SPRAY

(Contains 2 lbs. Methoxychlor and 2 lbs. Malathion per gallon)

ACTIVE INGREDIENTS:

Methoxychlor, Technical* 23.2%

Malathion (O,O-dimethyl dithiophosphate of
diethyl mercaptosuccinate) 23.2%

Aromatic Petroleum Derivatives 47.0%

INERT INGREDIENTS: 6.6%

Total 100%

*Equivalent to 20.42% 2,2-bis (p-methoxyphenyl) 1,1,1-tri-
chloroethane and 2.78% other isomers and reaction products.

CAUTION!

Keep Out of Reach of Children

See Side Panel for Additional Cautions

BACK RUBBING DEVICES FOR BEEF CATTLE ONLY: For the reduction of Horn Flies and lice dilute 2 gallons with 9 1/2 gallons Diesel fuel or furnace type oil. DO NOT use any other type of oil. Use one gallon of solution to saturate burlap on "back rubber;" soak burlap every 2 to 3 weeks. DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. Avoid breathing spray mist. Wash thoroughly after handling. Do not contaminate feed and foodstuffs. Do not store or use near fire or open flame.

This product should be used strictly in accordance with label directions and precautions.

This product is highly toxic to fish. Keep out of lakes, streams or ponds. Shrimp and crab will be killed at application rate recommended. Do not apply where these are important resources. Do not apply to water drainage areas where runoff or flooding will contaminate ponds, lakes or streams.

Malathion is a cholinesterase inhibitor. Consult veterinarian at first sign of adverse reaction.

Do not contaminate water by direct application, cleaning of equipment or disposal of waste. Apply only as specified on this label.

Do not re-use container. Destroy when empty.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of material not in strict accordance with directions.

7401-12

4/10/1972

Page 1 of 2

ACCEPTED

APR 10 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTER-
ED UNDER NO. 7901-71 SUBJECT
~~TO ATTACHED COMMENTS~~

272