

(Front Panel)

VPG

## RANGE CATTLE SPRAY

(Contains 2 lbs. Methoxychlor and 2 lbs. Malathion per gallon)

## ACTIVE INGREDIENTS:

Methoxychlor, Technical* . . . . .	23.2%
Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) . . . . .	23.2%
Aromatic Petroleum Derivatives . . . . .	47.0%
INERT INGREDIENTS: . . . . .	6.6%
Total . . . . .	100.0%

\*Equivalent to 20.42% 2,2-bis (p-methoxyphenyl) 1,1,1-tri-chloroethane and 2.78% other isomers and reaction products.

## CAUTION

KEEP OUT OF REACH OF CHILDREN

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS


NET CONTENTS      GALLONS

(Side Panel)

DIRECTIONS: For Use Against Certain Insects Infesting Range Cattle and Livestock.

PRECAUTIONS IN USING THIS PRODUCT: Do not treat animals under one month of age. When applying sprays, avoid contamination of feed, food containers and watering troughs. DO NOT APPLY TO DAIRY ANIMALS.

NOTE: Do not use this product as a dip treatment. Use only as a spray.



ANIMAL	PEST	RATE IN WATER	REMARKS
Beef Cattle (Range Cattle)	Lice	2 gal. per 100 gal. or 13 ozs. per 5 gal.	Treat animals thoroughly. Repeat application in 10 days and thereafter as needed.
	Ticks	2 2/3 gal. per 100 gal. or 17 ozs. per 5 gal.	Treat animals thoroughly. Repeat application at 2 week intervals as needed.
	Hornflies	2/3 gal. per 100 gal. or 13/20 ozs. per 5 gal.	Apply on back and neck of animals and repeat application at 10-14 day intervals.
Hogs	Lice	2 gal. per 100 gal.	Apply a complete coverage spray to animals (including the inside of ears), pens and litter. Repeat in 10 days and thereafter as needed. Sprayed hogs should be kept out of sun and wind for a few hours after treatment.
	Sarcoptic Mange	2 2/3 gal. per 100 gal. or 17 ozs. per 5 gal.	
Sheep & Goats	Lice Ticks Keds	2 2/3 gal. per 100 gal. or 17 ozs. per 5 gal.	Spray animals thoroughly. Repeat application after 2 or 3 weeks if necessary. DO NOT APPLY TO MILK GOATS.

44-5753

100000

BACK RUBBING DEVICES FOR BEEF CATTLE ONLY: For the reduction of Horn Flies and lice dilute 2 gallons with 9 1/2 gallons Diesel fuel or furnace type oil. DO NOT use any other type of oil. Use one gallon of solution to saturate burlap on "back rubber", soak burlap every 2 to 3 weeks. DO NOT MAKE ACCESSIBLE TO DAIRY ANIMALS.

CAUTION: Harmful if swallowed. Avoid contact with skin, eyes and clothing. Avoid breathing spray mist. Wash thoroughly after handling. Do not contaminate feed and foodstuffs. Do not store or use near fire or open flame.

This product should be used strictly in accordance with label directions and precautions.

This product is highly toxic to fish. Keep out of lakes, streams or ponds. Shrimp and crab will be killed at application rate recommended. Do not apply where these are important resources. Do not apply to water drainage areas where runoff or flooding will contaminate ponds, lakes or streams.

Malathion is a cholinesterase inhibitor. Consult veterinarian at first sign of adverse reaction.

Do not contaminate water by direct application, cleaning of equipment or disposal of waste. Apply only as specified on this label.

Do not re-use container. Destroy when empty.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

EPA Reg. No. 7401-121

44-3886-100

100-100