

Add 1 1/3 gallon of HI-YIELD DSMA LIQUID HERBICIDE to 40 gallons of water and apply to 1 acre. Apply as a post-emergence directed spray to grove floor in interspaces and around base of trees. Spray unwanted vegetation to just short of runoff. Do not exceed 3 applications per year. Do not allow spray to contact fruit, leaves, stems or bark of trees.


NON-CROP: HI-YIELD DSMA LIQUID HERBICIDE is useful for control of Johnsongrass, nutsedge, Dallisgrass, cocklebur, barnyardgrass, cogweed, sandbur, puncturevine and certain other weeds on drainage ditch banks, rights-of-way, fence rows, storage yards, and similar non-crop areas. Mix HI-YIELD DSMA LIQUID HERBICIDE at a rate of 1 to 2 gallons in 100 gallons of water. Spray unwanted vegetation at a rate of about 50 gallons of spray solution per acre. Additional surfactant is not necessary. Spray the unwanted vegetation thoroughly wetting plants to run off. Any spray equipment that gives good coverage may be used. If regrowth occurs, reapply as required.

DO NOT CONTAMINATE WATERS USED FOR DOMESTIC CONSUMPTION OR BY ANIMALS, WILDLIFE AND AQUATIC LIFE, OR IRRIGATION PURPOSES.

LAWN AND ORNAMENTAL TURF: HI-YIELD DSMA LIQUID HERBICIDE is useful for control of Dallisgrass, sandbur, bahiagrass, nutsedge, barnyardgrass, chickweed and wood sorrel with little or no injury to good lawn grasses. On new lawn, do not treat until after three mowings. Good grasses may be temporarily discolored. Zoysia, Bluegrasses and Bermuda are quite tolerant. Do not use on St. Augustine or centipede.

Mow lawns to 1 1/2 inches high before treatment. Mix 4 fluid ounces of HI-YIELD DSMA LIQUID HERBICIDE in 5 gallons of water and apply to an area of 1,000 square feet. Spray thoroughly to wet all undesirable plants. Repeat applications 10 to 14 days apart, may be needed for good control.

FBI Reg. No. 1-1-72



7401-140  
MP

(Front Panel)

Apply evenly to apply with fertilizer or seed

CONTAINS 40 LBS (18.18 kg)

Crabgrass

Year-Around

CRABGRASS AND WEED PREVENTION

- \* MAINTAINS CRABGRASS AND 26 OTHER ANNUAL WEEDS FROM  
THESE BROADLEAFED WEEDS IN LAWNS, BEDS AROUND PERENNIALS,  
FRUIT FLOWERS AND VEGETABLE GARDENS
- \* CAN BE EFFECTIVELY USED ON ST. AUGUSTINE, BERBERIS, PALM, PINE,  
DOGWOOD, MERION AND KENTUCKY BLUE GRASS, CARPET AND CYNODONT  
GRASS
- \* SPECIFICALLY RECOMMENDED FOR WEED CONTROL IN PERENNIALS,  
FRUIT FLOWERS, SHRUBS AND TREES
- \* MAY BE USED OVER UNDERLYING ROOTS OF DESIRABLE VEGETATION
- \* REMAINS EFFECTIVE UP TO 3 MONTHS
- \* CLEAN, FREE FLOWING AND ODORLESS
- \* IDEAL PRE-EMERGENCE WEED-KILLER TO INSURE WEED FREE AREAS

CAUTION (See Back)

KEEP OUT OF REACH OF CHILDREN

SEE BACK FOR DIRECTIONS


... ..  
... ..  
... ..

... ..  
... ..

NET WEIGHT 30 lbs.

(Back Panel)

Page 2.



NET WEIGHT 30 LBS.

ferti-lome

Year-Around

CRABGRASS AND WEED PREVENTER

4-9-0

GUARANTEED ANALYSIS:

4.00% Ammoniacal Nitrogen  
 12.00% Phosphoric Acid (P<sub>2</sub>O<sub>5</sub>)  
 12.00% Potash (K<sub>2</sub>O)

Contains Food Derived from: Sulfate of Ammonia  
 In all cases, "Minimum Guaranteed Analysis"

KEEP OUT OF REACH OF CHILDREN. Do not use on lawns.  
 Avoid contact with eyes. Wash after using.

Do not contaminate feed and foodstuffs. Do not use on  
 lawns, lawns, lawns or newly seeded lawns. Do not use on  
 lawns, lawns and other pesticides. This product is  
 used to avoid spilling chemical on lawn.

Do not reuse container. Destroy when empty.

ACTIVE INGREDIENT:

1,1,1-trichloro-2,2,2-trifluoroethane  
 1,1,1-trichloro-2,2,2-trifluoroethane

VOLUNTERS BUREAU

Bonham, Texas

ferti-lome

DIRECTIONS:

Dear Folks:

I recommend this package of ferti-lome Year-Around CRABGRASS AND  
 WEED PREVENTER because it is so effective and economical to use.