

ferti·lome®

ACCEPTED
Sep 7-1971
UNDER THE FEDERAL INSECTIONS
FUNGICIDE AND ROENTICIDE
FOR ECONOMIC POISON RESISTANCE
UNDER NO. 7471-135

DIRECTIONS

Dear Folks:

We feel that you will find the following directions easy to follow and this new product will aid in your lawn chinch bug control and lawn feeding programs.

ferti-lome CHINCH BUG CONTROL PLUS FERTILIZER is designed to perform two important functions: Control Chinch Bugs and supply a balanced, fast-acting fertilizer that will give you a greener lawn.

WHEN TO APPLY. Apply **ferti-lome CHINCH BUG CONTROL PLUS FERTILIZER** at the first sign of insects or insect damage. Since chinch bugs are most active during warm weather, they may start to show up in early spring. One treatment controls chinch bugs for at least 8 weeks. In some instances the lawn may not be troubled again for months. As a control against reinfestation follow the first treatment with a second application 8 weeks later.

HOW TO APPLY. For best results, treat entire lawn; do not spot treat. Apply the contents evenly to 3000 square feet (100' x 30') of lawn area. **AFTER APPLICATION, WATER LAWN THOROUGHLY TO WASH FERTILIZER DOWN INTO GRASS.**

The following spreader settings are calculated to deliver the recommended amount of product by covering the lawn in one direction. Do not use criss cross pattern. To insure complete coverage, overlap wheel tracks.

<u>SPREADER</u>	<u>ONE BAG COVERS 3,000 SQ. FT. SPREADER SETTING*</u>	To build good turf, feed with ferti-lome monthly during growing season
ferti-lome Cyclone	4½	
ferti-lome Deluxe	8	
Jackson	7	
Lawn Beauty	6	
Scott's	6½	
Seas	8	
Central	8	

2. HOW DO YOU USE IT?

WHEN TO APPLY. Apply **ferti-lome CHINCH BUG CONTROL PLUS FERTILIZER** at the first sign of insects or insect damage. Since chinch bugs are most active during warm weather, they may start to show up in early spring. One treatment controls chinch bugs for at least 8 weeks. In some instances the lawn may not be troubled again for months. As a control against reinfestation follow the first treatment with a second application 8 weeks later.

HOW TO APPLY. For best results, treat entire lawn; do not spot treat. Apply the contents evenly to 3000 square feet (100' x 30') of lawn area. **AFTER APPLICATION, WATER LAWN THOROUGHLY TO WASH FERTILIZER DOWN INTO GRASS.**

The following spreader settings are calculated to deliver the recommended amount of product by covering the lawn in one direction. Do not use criss cross pattern. To insure complete coverage, overlap wheel tracks.

<u>SPREADER</u>	<u>ONE BAG COVERS 3,000 SQ. FT. SPREADER SETTING*</u>	
ferti-lome Cyclone	4½	To build good turf, feed with ferti-lome monthly during growing season
ferti-lome Deluxe	8	
Jackson	7	
Lawn Beauty	6	
Scott's	6½	
Seas	8	
Central	8	

When used as directed, this new product will kill existing chinch bugs, control chinch bugs from reinfesting your lawns and feed your lawn in one application.

Sincerely,
the ferti-lome people


NOTICE: Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

*T. M. Reg. Stauffer Chemical Company for insecticide.

Net Wt. 40 Lbs.

ferti·lome®

CHINCH BUG CONTROL PLUS FERTILIZER 11-5-6



GUARANTEED FERTILIZER ANALYSIS, MIN.

Total Nitrogen (N).....	11.00%
10.26% Ammoniacal Nitrogen	
.74% Nitrate Nitrogen	
Available Phosphoric Acid (P ₂ O ₅)	5.00%
Water Soluble Potash (K ₂ O).....	6.00%
Derived from: Ammonium Sulfate, Ammoniated Superphosphate and Muriate of Potash.	
Soil Conditioners: Bone Meal, Cotton Seed Meal, Blood Meal, Rice Hulls.	
Ingredient Statement as an Insecticide	
ACTIVE INGREDIENT:	
O,O,O,O-Tetrapropyl dithiopyrophosphate †.....	1.32%
INERT INGREDIENTS:.....	98.68%
	Total..... 100.00%

†Equivalent 26.4 lbs. Aspon (O,O,O,O-Tetrapropyl dithiopyrophosphate) per ton.

READ ENTIRE LABEL USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

KEEP BAG CLOSED WHEN NOT IN USE.—STORE IN A DRY PLACE.

WARNING: May be fatal if swallowed. Avoid breathing dust. Do not get in eyes, on skin, on clothing. In case of con-

Top

corn, cottonseed meal, bone meal, cotton seed meal, blood meal, rice hulls.

Ingredient Statement as an Insecticide

ACTIVE INGREDIENT:

O,O,O,O-Tetrapropyl dithiopyrophosphate †..... 1.32%

INERT INGREDIENTS:..... 98.68%

Total..... 100.00%

†Equivalent 26.4 lbs. Aspon (O,O,O,O-Tetrapropyl dithiopyrophosphate) per ton.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, WARNINGS AND DIRECTIONS.

KEEP PESTICIDE IN ORIGINAL CONTAINER.

KEEP BAG CLOSED WHEN NOT IN USE.—STORE IN A DRY PLACE.

WARNING: May be fatal if swallowed. Avoid breathing dust. Do not get in eyes, on skin, on clothing. In case of contact, immediately flush eyes with plenty of water; for skin, wash with soap and water. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Keep children and pets off treated areas for 24 hours and until grass has completely dried. Dispose of container promptly when empty.

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

Fla. Reg. No. 370-1050

U.S.D.A. Reg. No. 7401-133

VOLUNTARY PURCHASING GROUPS, INC.

Bonham, Texas 75418

ferti·lome