

## IS THIS THE PACKAGE FOR?

This new pre-emergence weed-killer is for use on lawns and flower gardens. It kills Crabgrass, other annual weeds, and some perennial weeds. Use on lawns, around shrubs, and in flower gardens.

## HOW DO I USE IT?

**ON LAWNS AND PUTTING GREENS:** Use 23 pounds of product per acre. Apply material going one way then return and go the other way to insure necessary and complete coverage. Apply to the following weeds: CRABGRASS, ALBUQUERQUE, BERMUDA, BAHIA, ZONCH, KYLLIS, KENTUCKY BLUEGRASS, CARPET AND CENTIPEDE GRASS, ST. AUGUSTINE, FALLOUT, SANDSPUR (Sandspur), CARPET WELD, COMMON CHICKEN, FALLOUT, BERMUDA PANICUM, GREEN AND YELLOW FOXGLOVE, ORNAMENTAL GRASS, QUINQUARTER, PURSLANE, TEXAS MILLET, WITCHGRASS, MONARDRA, LOVE GRASS, SPOTTED AND PROSTRATE SPURGE, RED TOOTH, ANNUAL BLUE GRASS, STINKGRASS, REISIM, FALLOUT, MILLET, BARLEY, DODDER, BARNYARD GRASS, GOOSEGRASS, WILD RICE, CHERRY AND SASSANIA. The use of this product on putting greens should be restricted to portions of greens infested with crabgrass and bermuda which are known to be tolerant to Dacthal. Apply in early spring before crabgrass and annual weeds begin to germinate. If seeding is necessary, wait 30 days after application to a lawn. A single application should suffice for most green grasses. For annual bluegrass or late germinating grasses, a second application at a rate of 23 pounds of product per acre is recommended. Apply when grass leaves are dry. Water in after application. Do not rake lawn or disturb soil surface after application.


ONE BAG COVERS 2500 SQ. FT.

## SPREADER SETTINGS:

1	3
2	3 1/2
3	4
4	4 1/2
5	5
6	5 1/2
7	6
8	6 1/2
9	7
10	7 1/2

30 seconds use can only be taken as a guide. The rate of application has been thoroughly tested on new spreaders. The rate of application is affected by spreader, speed of pushing, roughness of lawn, etc. The operator should carefully observe rate of application and make the job and make corrective adjustments of spreader setting if necessary.

**IN FLOWER GARDENS:** Apply to soil on following list of specifically recommended flowers, shrubs and trees, in early spring after establishment or as supplemental application following establishment or setting out. Delayed or secondary applications can be made at any time provided beds have been cultivated and are weed free. Weed control up to 4 months or more may be expected following proper application:



$\frac{1}{2} \times 10 = 5$

774. 10

1002 N. 77th St.

1

2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

28

29

30

31

32

33

34

35

36

37

38

39

40

41

42

43

44

45

46

47

48

49

50

51

52

53

54

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

86

87

88

89

90

91

92

93

94

95

96

97

98

99

100

101

102

103

104

105

106

107

108

109

110

111

112

113

114

115

116

117

118

119

120

121

122

123

124

125

126

127

128

129

130

131

132

133

134

135

136

137

138

139

140

141

142

143

144

145

146

147

148

149

150

151

152

153

154

155

156

157

158

159

160

161

162

163

164

165

166

167

168

169

170

171

172

173

174

175

176

177

178

179

180

181

182

183

184

185

186

187

188

189

190

191

192

193

194

195

196

197

198

199

200

201

202

203

204

205

206

207

208

209

210

211

212

213

214

215

216

217

218

219

220

221

222

223

224

225

226

227

228

229

230

231

232

233

234

235

236

237

238

239

240

241

242

243

244

245

246

247

248

249

250

251

252

253

254

255

256

257

258

259

260

261

262

263

264

265

266

267

268

269

270

271

272

273

274

275

276

277

278

279

280

281

282

283

284

285

286

287

288

289

290

291

292

293

294

295

296

297

298

299

300

301

302

303

304

305

306

307

308

309

310

311

312

313

314

315

316

317

318

319

320

321

322

323

324

325

326

327

328

329

330

331

332

333

334

335

336

337

338

339

340

341

342

343

344

345

346

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

405

406

407

408

409

410

411

412

413

414

415

416

417

418

419

420

421

422

423

424

425

426

427

428

429

430

431

432

433

434

435

436

437

438

439

440

441

442

443

444

445

446

447

448

449

450

451

452

453

454

455

456

457

458

459

460

461

462

463

464

465

466

467

468

469

470

471

472

473

474

475

476

477

478

479

480

481

482

483

484

485

486

487

488

489

490

491

492

493

494

495

496

497

498

499

500

501

502

503

504

505

506

507

508

509

510

511

512

513

514

515

516

517

518

519

520

521

522

523

524

525

52

Figure 1. The effect of the concentration of the *Agrobacterium* suspension on the transformation efficiency of *Agrobacterium* strains. The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D). The concentration of the *Agrobacterium* suspension was 10<sup>6</sup> cells/ml (A), 10<sup>7</sup> cells/ml (B), 10<sup>8</sup> cells/ml (C), and 10<sup>9</sup> cells/ml (D).

100

[illegible]

\_\_\_\_\_

9

\_\_\_\_\_

\_\_\_\_\_

\_\_\_\_\_

need control up to 6 months or more may be expected following broadcast application. See above list for specifically recommended list of flowers, shrubs and trees.

... and other seeds which remain in the soil.

126

**NON-WARRANTY:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

1. WHAT IS THIS PACKAGE FOR?

ferti-lome SPOT WEED KILLER is formulated for "spot" applications to undesirable broadleaved weeds in your lawns.

WEEDS THIS PRODUCT CONTROLS

Dandelion, Poison Ivy, Plantain, Chickweed, Wild Onion, Ground Ivy, Pennywort, Canadian Thistle, Common and Giant Ragweeds, Bindweed (Annual Morning Glory), Wild Garlic, Stinging Nettle, Poison Hemlock and Redroot Pigweed.

2. HOW DO YOU USE IT?

SHAKE WELL BEFORE USING. Remove protective cap. Stand close to weed (1 to 2 feet). Point arrow on nozzle at weed to be killed. Press Nozzle to spray center of weed. Spray about 1 second. One application will kill most weeds listed. If certain tough weeds are not dead within 3 weeks, repeat application.

NOTE: Be careful when treating weeds not to spray surrounding desirable grasses unless the browning of these grasses can be tolerated. Also avoid spraying or splattering of nearby flowers and other desirable plants.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

This new product, used according to directions, will control listed undesirable weeds in your lawn.

**CAUTION:** Avoid contact with skin, eyes and clothing. Wash hands after use. Do not store near fertilizers, seeds, insecticides or fungicides.

**WARNING:** Contents under pressure. Do not puncture. Exposure to high temperature may cause bursting. Keep at room temperature—away from direct sunlight, radiators, stoves, hot water and other heat. Never throw container into fire or incinerator.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Active Ingredient:

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	0.89%
Inert Ingredients:	99.11%

Total 100.00%

\*Equivalent to 0.74% 2,4-Dichlorophenoxyacetic acid

MANUFACTURED FOR VOLUNTARY PURCHASING GROUPS, INC., BONHAM, TEXAS

U.S.D.A. Reg. No. 7401-144

14M-1-71



ferti·lome



New SPOT WEED KILLER

- CONTROLS DANDELION, POISON IVY, PLANTAIN, CHICKWEED, WILD ONION, GROUND IVY, PENNYWORT AND CERTAIN OTHER LISTED BROADLEAVED WEEDS
- IDEAL FOR SPOT TREATMENT IN LAWNS
- FAST, EASY AND EFFECTIVE
- CONTAINS 2,4-D . . . A HORMONE TYPE WEED KILLER



NET WEIGHT ONE POUND (16 OUNCES)

**CAUTION:**  
KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

ACCEPTED WITH COMMENTS

ACCEPTED

8/9/71

# 1. WHAT THE PACKAGE TOLD YOU

ferti-lome SPOT WEED KILLER is formulated for "spot" applications to undesirable broadleaved weeds in your lawns

## WEEDS THIS PRODUCT CONTROLS

Dandelion, Poison Ivy, Plantain, Chickweed, Wild Onion, Ground Ivy, Pennywort, Canadian Thistle, Common and Giant Ragweeds, Bindweed (Annual Morning Glory), Wild Garlic, Stinging Nettle, Poison Hemlock and Redroot Pigweed.

## 2. HOW DO YOU USE IT?

**SHAKE WELL BEFORE USING.** Remove protective cap. Stand close to weed (1 to 2 feet). Point arrow on nozzle at weed to be killed. Press Nozzle to spray center of weed. Spray about 1 second. One application will kill most weeds listed. If certain tough weeds are not dead within 3 weeks, repeat application.

**NOTE:** Be careful when treating weeds not to spray surrounding desirable grasses unless the browning of these grasses can be tolerated. Also avoid spraying or splattering of nearby flowers and other desirable plants.

## 3. HERE ARE THE RESULTS YOU MAY EXPECT:

This new product, used according to directions, will control listed undesirable weeds in your lawn.

**CAUTION:** Avoid contact with skin, eyes and clothing. Wash hands after use. Do not store near fertilizers, seeds, insecticides or fungicides.

**WARNING:** Contents under pressure. Do not puncture. Exposure to high temperature may cause bursting. Keep at room temperature, away from direct sunlight, radiators, stoves, hot water and other heat. Never throw container into fire or incinerator.

**NOTICE:** Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

### Active Ingredient

*Dimethylamine salt of 2,4-Dichlorophenoxyacetic acid	0.89%
Inert Ingredients	99.11%

Total	100.00%
-------	---------

\*Equivalent to 0.74% 2,4-Dichlorophenoxyacetic acid

**MANUFACTURED FOR VOLUNTARY PURCHASING GROUPS, INC., BONHAM, TEXAS**

U.S.D.A. Reg. No. 7401-144

14M-1-71



ferti-lome



# New SPOT WEED KILLER \$1.98

- CONTROLS DANDELION, POISON IVY, PLANTAIN, CHICKWEED, WILD ONION, GROUND IVY, PENNYWORT AND CERTAIN OTHER LISTED BROADLEAVED WEEDS
- IDEAL FOR SPOT TREATMENT IN LAWNS
- FAST, EASY AND EFFECTIVE
- CONTAINS 2,4-D... A HORMONE TYPE WEED KILLER



NET WEIGHT ONE POUND (16 OUNCES)

**CAUTION:**  
KEEP OUT OF REACH OF CHILDREN  
SEE BACK PANEL FOR ADDITIONAL CAUTIONS