

CAUTION:

Keep out of reach of children.
See label for additional cautions.

10% SEVIN* DUST

ACTIVE INGREDIENTS:

Carbaryl (1-naphthyl N-methylcarbamate)

10%

INERT INGREDIENTS:

90%

*Trade-Mark of Union Carbide Corporation for the active ingredient, 1-naphthyl N-methylcarbamate.

100%

CAUTION: HARMFUL IF INHALED OR SWALLOWED. Do not breathe dust. Do not take internally. Avoid breathing of dust. Skin contact may be harmful. Avoid contact. Wash hands before eating. Take shower or bath after work. Wear regular long sleeved work clothing. Change to clean clothing daily. Note for Physician: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

Do not reuse container for any purpose. Destroy empty container.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS.

DIRECTIONS FOR USE — Vegetable, Vine and Field Crops

Sevin insecticide will not control insects if used with lime or other highly alkaline materials. Apply when air is calm, using ground equipment, aircraft or hand duster. Unless otherwise specified, apply at first sign of infestation and repeat at 7-10 day intervals when needed to maintain control. All dosages are given in pounds per acre. Observe use limits given for specific crops.

ALFALFA, CLOVERS, COWPEAS, PEANUTS AND SOYBEANS—Use 10 lbs. for blister beetles and Mexican bean beetle. Use 10 lbs. for alfalfa caterpillar, bean leaf beetle, cucumber beetles, green clover worm, Japanese beetle, leafhoppers, three-cornered alfalfa hopper, thrips and velvetbean caterpillar. For armyworms, corn earworm, stink bugs and webworms, use 10-15 lbs. Some leaf injury may occur if applications are made when tender foliage is wet. May be applied up to and including the day of harvest.

BEANS—Use 10 lbs. for Mexican bean beetle; use 10 lbs. for bean leaf beetle, cucumber beetles, flea beetles, Japanese beetle, leafhoppers, velvetbean caterpillar and western bean cutworm. Use 10-15 lbs. for armyworms and tarnished plant bugs. 20 lbs. for corn earworms, lima bean pod borer, lygus and stink bugs. May be applied up to and including the day of harvest.

CUCUMBERS, MELONS, PUMPKIN AND SQUASH—Use 10 lbs. for pickleworm and melonworm; 10 lbs. for cucumber beetles, flea beetles, leafhoppers and

squash bugs. Avoid excessive applications. Some leaf injury may occur when tender foliage is wet or in the presence of prolonged humidity. May be applied up to and including the day of harvest.

COTTON—Early season control of: Cotton leafworm, cotton fleahopper, flea beetles, striped blister beetle, thrips. Apply 10 lbs. per acre. Mid and late-season control of: Boll weevil, bollworm, fall armyworm, cotton leaf perforator, leaf rollers, leafhoppers, lygus bugs (light to moderate infestations in Western irrigated cotton), tarnished plant bugs, aphids (populations will be suppressed by repeated applications of this insecticide).—Apply 10-20 lbs. For pink bollworm—apply 15-25 lbs. on a 5-7 day schedule depending on the size of the cotton and degree of infestation. For salt marsh caterpillar, stink bugs—apply 20 lbs. For grasshoppers—apply 10-15 lbs. The higher dosage should be used when grasshoppers are mature or material is applied on partly defoliated plants. Allow 7 days before grazing or harvesting crop residue for feed or bedding for meat and dairy animals.

POTATO, TOMATO, EGGPLANT AND PEPPER—Use 10 lbs. for Colorado potato beetle, flea beetles, and leafhoppers. Use 10-20 lbs. for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. May be applied up to and including the day of harvest.

OBSERVE THE FOLLOWING PRECAUTIONS

Toxic to honey bees. Do not apply during bloom when honey bees or other pollinators are active. When necessary to use during such periods, warn beekeepers well in advance to locate hives at a safe distance until one week after application. Avoid drift to adjoining food and forage crops.

The Federal Food and Drug Administration has set official tolerances for certain pesticide residues. In order that residues on food and forage crops will not exceed established tolerances, use only at recommended rates, and do not apply closer to harvest than specified. Consult local agricultural authorities for additional information, as the timing, number, and rate of applications needed will vary with local conditions.

NON-WARRANTY: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS

NET CONTENTS 50 LBS.
2M-1-22

EPA Reg. No. 7401-154

ACCEPTED