

# TOMATO and VEGETABLE DUST

## 1. WHAT IS THIS PACKAGE FOR?

This product is especially formulated to control listed insects and plant diseases in your tomato and vegetable garden.

## 2. HOW DO YOU USE IT?

Use the squeeze duster to unpack contents that may have settled during shipment. **Remove protective cap by cutting flap.** For best results hold at 45° angle from ground. Hold squeeze duster 12-18 inches from surfaces to be treated. Squeeze duster and apply dust in such a manner that a uniform, hardly visible coating results on both sides of foliage, stems and all above ground parts. Use at the rate of 1 lb. per 1200 sq. ft. except on cucumbers, squash, melons and peas use at the rate of 1 lb. per 2000 sq. ft. Start application before insects and/or diseases appear. Repeat at 7-10 day intervals throughout growing season. Replace protective cap.

**TOMATOES:** *blight* Heliothis, Fruitworm, Flea Beetle, Early and Late Blight, Anthracnose, *Septoria Leaf Spot*, *Septoria Leaf Spot*, Flea Beetle, European Flea Beetle, Tarnished Plant Bug, Colorado Potato Beetle, Anthracnose, Peronospora on peppers, *Phomopsis*, *Alternaria* and *Ascochyta* on eggplant.

**CUCUMBERS, SQUASH AND MELONS:** Striped and Spotted Cucumber Beetle, Squash Bug, Flea Beetle, Anthracnose, and Angular Leaf Spot and Scab. Application may be made on above vegetables up to day of harvest. *Mexican Bean Weevil*, *Bean Leaf Beetle*, Mexican Bean, Bean Leaf Flea, and Japanese Beetles, Leafhopper, *Downy Mildew*, *Rust* and *Anthracnose*.

**BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, BROCCOLI AND KOHLRAB:** Cabbage Worm, Flea Beetle, Harlequin Bug, Corn Earworm, Armyworm, Cabbage Caterpillar.

**BEANS AND PEAS:** Leafhopper and Pea Weevil, respectively. Do not apply to beans, brussels sprouts, cabbage, cauliflower, broccoli or kohlrabi within 7 days of harvest. Do not apply to peas within 10 days of harvest. Do not apply to kohlrabi after edible parts are half grown.

Use on **CORN:** European Corn Borer, dust every 5-7 days from the time eggs are first laid (when corn is knee high on most varieties). Corn Earworm, dust when insects are first seen. Make at least 3 applications at 2-3 day intervals. Do not apply this product to the tassel region of corn during the pollen shed period to seriously reduce bee population.

Do not apply to corn up to day of harvest. Do not feed forage to dairy animals or animals being finished for slaughter.

## 3. HERE ARE THE RESULTS YOU MAY EXPECT!


When TOMATO & VEGETABLE DUST, used as directed will kill listed insects and control listed diseases that ruin tomatoes and vegetables, thus insuring a more bounteous harvest.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**




ferti-lome



# TOMATO and VEGETABLE DUST

\$1.89



740-159  
**ACCEPTED**

OCT 18 1973

UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR REGISTERED USE SUBJECT TO ATTACHED COMMENTS.

ACTIVE INGREDIENTS

...  
...  
...  
...  
INERT INGREDIENTS

BEANS: Mexican Bean, Bean Leaf, Flea, and Japanese Beetles, Leafhopper, Edema, Downy Mildew, Rust and Anthracnose

BRUSSELS SPROUTS, CABBAGE, CAULIFLOWER, BROCCOLI AND KOHLRABBI: Cabbage Worm, Flea Beetle, Harlequin Bug, Corn Earworm, Armyworm, Cabbage Caterpillar

PEAS AND PEAS: Leafhopper and Pea Weevil, respectively. Do not apply to beans, brussels sprouts, cabbage, cauliflower, broccoli or peas within 7 days of harvest. Do not apply to peas within 10 days of harvest. Do not apply to kohlrabi after edible parts are half grown.

WAX: CORN: European Corn Borer, dust every 5-7 days from the time eggs are first noticed (when corn is knee high on most varieties). Corn Earworm, dust when silks appear. Make at least 3 applications at 2-3 day intervals. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

NOTE: May be applied to corn up to day of harvest. Do not feed forage to dairy animals or animals being finished for slaughter.

**3. HERE ARE THE RESULTS YOU MAY EXPECT!**

For some TOMATO & VEGETABLE DUST, used as directed will kill listed insect and control listed diseases that ruin tomatoes and vegetables, thus assisting in producing a more bounteous harvest.

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**

May be fatal if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes and clothing. Do not breathe dust. Wash thoroughly after using.

TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal.

This product is toxic to fish and wildlife. Birds feeding on treated area may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from area treated. Apply only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do not reuse container for any purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.


EPA Reg. No. 7401-159

**VOLUNTARY PURCHASING GROUPS, INC.**  
BONHAM, TEXAS 75418

\*Sevin, Reg. Trademark of Union Carbide Corp. for Carbaryl Insecticide.

*Bean leaf beetle*  
**7401-159**

**ACCEPTED**  
**OCT 18 1975**  
UNDER THE FEDERAL INSECTICIDE, FUNGICIDE AND RODENTICIDE ACT FOR REGISTERED USE. SUBJECT TO ATTACHED COMMENTS.



**ACTIVE INGREDIENTS:**  
Carbaryl (1-Naphthyl N-methylcarbamate) 90%  
Zineb (Zinc Ethylene Bisdithiocarbamate) zinc content 10%

**INERT INGREDIENTS:**  
Fineness: Not less than 100 mesh

**KILLS HORNWORMS, ARMYWORMS, BEETLES, LEAF HOPPERS AND OTHER INSECTS AND CONTROLS DOWNY MILDEW, ANTHRACNOSE, GREY LEAF SPOT AND OTHER DISEASES ON TOMATOES AND VEGETABLES.**

**CONTAINS EFFECTIVE INSECTICIDE**

**NET WEIGHT 10 OUNCES**

**CAUTION**

**KEEP OUT OF REACH OF CHILDREN**  
**SEE BACK PANEL FOR ADDITIONAL CAUTIONS**

