

Best Available Copy

4444
2222

7401-163

10/06/1972

1/1

1. WHAT IS THIS PACKAGE FOR?
This spray gives the Azalea, Camellia and Gardenia grower a spray which solves many insect and disease problems.

2. HOW DO YOU USE IT?
Carefully follow directions in table below:

INSECT	DOSAGE	REMARKS
Aphids, Mites, Lace Bugs, Mealybugs, Whiteflies, Thrips	2 tablespoons per gallon of dilute spray	Apply thoroughly to all foliage surfaces when insects or their damage first appear. Repeat at 10-14 day intervals as necessary to maintain control. For Florida red scale apply when crawlers have settled on foliage.
Azalea Scale, Florida red scale, Fletcher scale, Tea Scale, Soft Scale	4 tablespoons per gallon of dilute spray	Apply to 25 sq. ft. Spray ground and surrounding area under plants. Begin 4 weeks before bloom. Repeat at 3-4 week intervals through bloom period.

DISEASE	DOSAGE	REMARKS
Azalea petal blight (Ovulinia), Camellia flower or petal blight (Sclerotinia)	Mix 1 pint in 10 gallons of water	Apply to 25 sq. ft. Spray ground and surrounding area under plants. Begin 4 weeks before bloom. Repeat at 3-4 week intervals through bloom period.

3. HERE ARE THE RESULTS YOU MAY EXPECT!
When used as directed, this new formulation controls the listed damaging insects and disease on Azaleas, Camellias and Gardenias.

CAUTION: May be harmful if swallowed, inhaled or absorbed through the skin. May cause irritation of the eyes, nose, throat and skin. Do not take internally. Do not breathe fumes or spray mist. Avoid contact with skin, eyes and clothing. Wash thoroughly after handling. Avoid contamination of food and food-stuffs.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.


Food utensils such as a teaspoon or measuring cup should not be used for food purposes after use with insecticides.

Do not use container for any purpose. Destroy empty container. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.


VOLUNTARY PURCHASING GROUPS, INC.
BONHAM, TEXAS

28M-1-102 EPA Reg. No. 7401-163

NOTE: Do not use on ferns or Junipers.



ferti-lome



AZALEA-CAMELLIA GARDENIA PROBLEM SOLVER

- KILLS APHIDS, MITES, LACE BUGS, WHITE FLY, MEALY BUGS, LISTED SCALE INSECTS AND CONTROLS BLOSSOM BLIGHT
- CONTAINS INSECTICIDE-MITICIDE-SCALICIDE-FUNGICIDE


Active Ingredients:

Moluthion	7.5%
Fenitrothionbenzene	12.5%
Inert Ingredients	80.0%

*0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

NET CONTENTS 16 FL. OZS. (ONE PINT)

**CAUTION: KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS**



ACCEPTED

 OCT 6 1972

 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER-
 ED UNDER NO. 7401-163