- July 21/1972

ikerisin llorogie

Liquid \$198
NUTGRASS KILLER

- Ready to use no nucley necessary Treats up to 4000 nurgess; plants
 Contains Sprunder Studen for more effective control.

NET CONTENTS 16 FL. OZ. (ONE PINT)

CAUTION

Keep Out of Reach of Children See Back Panel for Additional Cautions

Voluntary Purchasing Graups, Inc.

BRSALIBE

(分形) ATT サーノ・イープログ でACCOVAL FRONT G LICK | BOH | ではア COLOR あたっただと

I WHAT IS THIS PACKAGE FOR?

The second of th

2. HOW DO YOU USE 17?

10'S DOD SO WELL SESSION AND ALL WOLF THE FILE AL IMPECTIONE
21 OF THE SOLUTION AT THE SOLUTION AND FOR FORECHINANCE TO SENTICIPE ANY
plants may propose a papear from nots the ground Sente applications during the
ground Sente inay the necessary to cean of

ACCEPTED July - 27-1973.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

15-1-36 are ted this product wais indicate - Ifaces 3-3-5 (1-22 for presing of nuture), parts simetimen causes nuturass to multiply

plants simetimal causes nutgrass to noutgy to be a compared to the condition of same and a compared to the com

Do not apply to desire: elegitation

Commit containmenter water's used for domestic consumption of the sometic weighter or advance of an applicable of the national section of the sometic constraints and call a privile on an account of the national section of

Do not to se Untainer for any purpose Destroy when empty

th wer assumes an insks of use, storage or handing of this mater is, our instruct accordance with directions given herewith

EPA Reg No 7401-168

SN 25 2 14

POTAL FRONT MICHA SOIL. Million Court

1. WHAT IS THIS PACKAGE FOR?

ferti-lome Systemic WEED KILLER is especially formulated to control weeds in lawns, golf courses and similar turf areas. See side of container for a partial list of weeds controlled.

2. HOW DO YOU USE IT?

Mix 10 tablespoonfuls in 6 gallons of water. Apply thoroughly to 1000 sq. ft. ($10' \times 100'$). Heavily infested lawns may require repeat application to kill missed weeds. Best results are obtained when weeds are young and activly growing. Application in early Spring and Fall when moisture and temperature make weeds grow rapidly is a good time to spray. Watering and fertilizing schedules should be maintained before and after spraying. Do not apply during periods of drought. If hose and sprayer is used, follow manufacturer's directions for proper mixing. Coarse sprays are less likely to drift. Do not apply within the dripline of desirable trees and shrubs.

For use in St. Augustine Grass mix 5 tablespoonfuls in 12 gals, of water. Apply thoroughly to 2000 sq. ft. of lawn area. Repeat application in 10-14 days if necessary. NOTE: May cause injury to St. Augustine Grass.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

ferti-lome Systemic WEED KILLER provides an economical and highly effective means of controlling weeds in your lawn thus letting your lawn grasses grow vigorously to provide you with a beautiful weed free lawn.

CAUTION: Harmful if swallowed or absorbed by skin. Causes irritation of eyes and skin. Do not get in eyes. Do not allow contact with skin and clothing. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention. Wash exposed skin with plenty of soap and water.

Direct spray at undesirable plants. Do not use the same spray equipment for applying other material to plants, as injury will result. Do not store near fertilizer, seeds, insecticides or fungicides. Avoid contamination of irrigation ditches and water used for domestic purposes. If lawn begins to brown temporarily, water heavily.

Do not use on Centipede grass, Dichondra, nor on lawns of turf where desirable Bent-grass or Clovers are present. Do not use on newly seeded lawns or turf.

Avoid spray drift to susceptible plants, as this product may injure cotton, beans, peas, grapes, ornamentals, etc.


Do not reuse container for any purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

28M-1-62

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS

EPA Reg. No. 7401-171



T. WHAT IS THIS PACKAGE FOR?

ferti-lome Systemic WEED KILLER is especially formulated to control weeds in lawns, golf courses and similar turf areas. See side of container for a partial list of weeds controlled.

2. HOW DO YOU USE IT?

Mix 10 tablespoonfuls in 6 gallons of water. Apply thoroughly to 1000 sq. ft. (10' x 100'). Heavily infested lawns may require repeat application to kill missed weeds. Best results are obtained when weeds are young and activly growing. Application in early Spring and Fall when moisture and temperature make weeds grow rapidly is a good time to spray. Watering and fertilizing schedules should be maintained before and after spraying. Do not apply during periods of drought. If hose and sprayer is used, follow manufacturer's directions for proper mixing. Coarse sprays are less likely to drift. Do not apply within the dripline of desirable trees and shrubs.

For use in St. Augustine Grass mix 5 tablespoonfuls in 12 gals, of water. Apply thoroughly to 2000 sq. ft. of lawn area. Repeat application in 10-14 days if necessary. NOTE: May cause injury to St. Augustine Grass.

3. HERE ARE THE RESULTS YOU MAY EXPECT!

ferti-lome Systemic WEED KILLER provides an economical and highly effective means of controlling weeds in your lawn thus letting your lawn grasses grow vigorously to provide you with a beautiful whed free lawn.

CAUTION: Harmful if swallowed or absorbed by skin. Causes irritation of eyes and skin. Do not get in eyes. Do not allow contact with skin and clothing. In case of contact, flush eyes with plenty of water for at least 15 minutes and get medical attention. Wash exposed skin with plenty of soap and water

Direct spray at undesirable plants. Do not use the same spray equipment for applying other material to plants, as injury will result. Do not store near fertilizer, seeds, insecticides or fungicides. Avoid contamination of irrigation ditches and water used for domestic purposes. If lawn begins to brown temporarily, water heavily.

Do not use on Centipede grass, Dichondra, nor on lawns of turf where desirable Bent-grass or Clovers are present. Do not use on newly seeded lawns or turf

Avoid spray drift to susceptible plants, as this product may injure cotton, beans, peas, grapes, ornamentals, etc.

Do not reuse container for any purpose. Destroy empty container.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

28M-1-62

VOLUNTARY PURCHASING GROUPS, INC. BONHAM, TEXAS

EPA Reg. No 7401-171



ferti·lome





- CONTROLS BROADLEAF WEEDS AND VINING TYPE WEEDS BY SYSTEMIC AND CONTACT ACTION
- 10 TABLESPOONS MAKE 6 GALLONS OF SPRAY THAT WILL COVER 1000 SQ. FT.

ACTIVE INGREDIENTS Dimethylamine Salt of Dicamba

(3,6-dichloro-o-anisic acid)*

Dimethylamine Salt of related acids

Dimethylamine Salt of 2,4-Dichloro-

*Equivalent to 1.19% 3,6-dichloro-o-anisic acid (Dicamba) **Equivalent to 6.84% 2,4-dichlorophenoxyacetic acid

NET CONTENTS 16 FL. OZS. (ONE PINT)

CAUTION, KEEP OUT OF REACH OF CHILDREN, SEE BACK PANEL FOR ADDITIONAL CAUTIONS

CONTROLS

Dandelion Clovers Chicory Docks Dog Fennel English Daisy Frenchweed Erect Knotweed Ground Ivy Hawkweed Healall Lambsquarter Mallow Morning Glory Henbit Oxalis Mustard Pennywart Peppergrass Pigwe**c**d Plantain (Buckhorn, Broadleaf, Rugel st Poison Oak Prostrate Knotweed Purslane Ragweed Sheep Sorrel Mouseear Chickweed Spotted Spurge Stitch Wort Wild Aster Wild Carrot Wild Garlic Wild Onion W.ld Radish Wood Sorrel Yarrow Puncturevine Milkweed Bull Thistle Shepherd's Purse

