

FH 19


7401-172

1044

Front Panel

ferti-lome

Fruit Tree Spray



* Provides Insecticide - Miticide - Scalecide for Control of Aphid, Codling Moth, Pear Psylla, Japanese Beetle on Apple, Pear, Peach, Apricot, Cherry, Nectarine and Quince.

* 3 Tablespoons Make 1 Gallon of Spray

ACTIVE INGREDIENTS:

) Methoxychlor Technical*..... 15%

Malathion**..... 12%

INERT INGREDIENTS***..... 73%

* Equivalent to 13.2% 2,2 bis(p-methoxyphenyl)-1,1,1-trichloroethane and 1.8% other isomers and reaction products.

** 0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate.

*** Contains Aromatic Petroleum Solvents.

EPA. Reg. No. 7401-172

EPA EST.NO. 7401-TX-1

KEEP OUT OF REACH OF CHILDREN

) CAUTION

SEE BACK PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

2044

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABEL DIRECTIONS.

DIRECTIONS FOR USE. READ ENTIRE LABEL DIRECTIONS.

1. What is this package for?

Fast, sure insect control is provided for your fruit trees when using ferti-lome Fruit Tree Spray.

2. HOW DO YOU USE IT?

Mix required dosage in water as listed in table below. Apply as a thorough cover spray to all surfaces of foliage and fruit. Repeat in 10-14 day if necessary to maintain control. Use only as specified on this label.

NOTE tbs=tablespoon, tsp=teaspoon

The following application schedule for the orchard pests; Codling Moth, Apple Maggot, Oriental Fruit Moth and Plum Curculio must be followed: Apply at petal fall and every 10 to 14 days thereafter until control is achieved.

CROP	INSECTS	RATES	REMARKS
Apples, Pears	Red Banded Leaf Roller, Codling Moth, Plum Cur- culio.	3 tbs. per gal. or 1 qt. in 21 gals.	May injure McIntyre and Cortland varieties. Do not apply to apples and pears within 7 days of harvest.
	Tent Caterpillar and Bud Moth (apples only) Mealy Bug, Pear Psylla, Japanese Beetle, Leafhopper, European and Two Spotted Mites	5 tsp per gal. or 1 qt. in 42 gals.	
Peaches	Plum Curculio, Oriental Fruit Moth, Terripan and Cottony Peach Scale.	3 tbs per gal. or 1 qt in 21 gals.	Do not apply to peaches within 21 days of harvest.
	Codling Moth, Aphids, Apple Maggot, Jap- anese Beetle, Leaf- hoppers, European Red and Two Spotted Mites.	5 tsp per gal. or 1 qt. in 42 gals.	
Apricots Cherries Nectarines Quinces	Plum Curculio, Codling Moth, Oriental Fruit Moth	3 tgs. per gal. or 1 qt. in 21 gals.	Do not apply to Apricots, Nectarines within 21 days of harvest. Do not apply to Cherries within 12 days of harvest or Quinces within 7 days Care should be used with Cherry since injury may result. May cause spotting on nectarines.
	Codling Moth, Spider Mites, Oriental Fruit Moth, Aphids, Cherry Fruit worm, Cherry Fruit Fly, Japanese Beetle and Leafhoppers	5 tsp per gal. or 1 qt in 42 gals.	

3. HERE ARE THE RESULTS YOU MAY EXPECT!

- Outstanding, easy and economical control of listed insects may be had on your valuable fruit trees with the use of ferti-lome Fruit Tree Spray

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

2/10/41

AND WHO

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH SKIN

Avoid breathing spray mist. Avoid contact with skin, eyes, or clothing. May be a skin sensitizer. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

This product is toxic to fish and aquatic invertebrates. Do not apply directly to water and wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.

This product is toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow to drift to blooming crops or weeds if bees are visiting the treatment area.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IN INHALED: Remove victim to fresh air. Give artificial respiration if indicated.

IF ON SKIN: Wash with plenty of soap and water.

IN IN EYES: Flush with plenty of water. Get medical attention if irritation persists.

NOTE TO PHYSICIAN: This product may cause cholinesterase inhibition. Atropine is antidotal. 2-PAM may be effective as an adjunct to atropine.

Voluntary Purchasing Group warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the Directions For Use. VPG makes no other expressed or implied warranty, including any other expressed or implied warranty, of FITNESS or of MERCHANTABILITY. In no event shall VPG be liable for indirect or consequential damages. This warranty shall not be changed by oral or written agreement unless signed by a duly officer of.

Voluntary Purchasing Groups, Inc.

Bonham, Texas 75418

NET CONTENTS

FL.OZS.