



5169

DUSTING SULFUR

- CONTROLS BLACK SPOT, POWDERY MILDEW, LEAF SPOT, RUST AND BROWN CANCKER ON ROSES
- CONTROLS CERTAIN FUNGI AND CERTAIN INSECTS ON LISTED ORNAMENTALS, FLOWERS, SHRUBS, FRUIT TREES AND VEGETABLES
- CONTROLS THRIPS, RUST MITES, RED SPIDER MITES AND TWO SPOTTED MITES ON CITRUS
- CONTROLS CHIGGERS
- MAY BE APPLIED AS DUST OR SPRAY



Active Ingredient:	
Sulfur	94%
Inert Ingredients:	6%
Total	100%

NET WEIGHT 32 OZ. (TWO POUNDS)

CAUTION KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR ADDITIONAL CAUTIONS

1. WHAT IS THIS PACKAGE FOR?

Produced for efficient, economic control of listed plant diseases and insects. May be used as a dust or, when suspended in water, as a convenient spray.

2. HOW DO YOU USE IT?

When used as a dust, punch out holes in top of shaker can and apply to all leaf surfaces for thorough coverage.

When used as a water spray, mix 4 tablespoons in 1 gallon of water. Spray leaf surfaces for thorough coverage. All dosages given are for dilute sprays. Spray often enough to keep new growth covered.

ROSES, ORNAMENTALS, SHRUBS, FLOWERS. (As Fungicide and Insecticide) for Black Spot, Brown Canker, Leaf Spot, Powdery Mildew, Thrips and Mites on Roses. Begin spraying when new growth starts in spring. Repeat at 7 day intervals and after rains. For disease control on ornamentals, shrubs and flowers begin application when diseases appear and repeat at 5-10 day intervals and after rains.

For control of Powdery Mildew, Thrips and Mites on **Azalea, Alder, Buttonbush, Catalpa, Cosmos, Crape Myrtle, Dahlia, Daisy, Delphinium, Aster, Calendula, Dogwood, Euonymus, Gladiolus, Golden Fleece, Hibiscus, Holly, Hydrangeas, Lady's Mantel, Ligustrum, Lilac, Petunia, Poplars, Silver Vine, Spirea, Sunflower, Sweet Pea, Verbena, Violets, Willows, Carnations, Cedars, Cherry Laurel, Chrysanthemums, Hollyhock, Juniper, Simlax and Spruce;** for control of Leaf Spot, Thrips and Mites on **Calendula, Cedars, Cherry Laurel, Chrysanthemum, Clematis, Columbine, Foxglove, Hollyhock, English Ivy, Juniper, Laurel, Sage, Simlax, Snapdragons and Spruce;** for control of Rust, Thrips and Mites on **Aster, Carnations, Chrysanthemums, Hollyhock and Snapdragons.**

VEGETABLES: (As Fungicide)

For control of Powdery Mildew on **Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower and Peas;** for control of Rust on **Beans and Peas.** Begin applications at first sign of disease. Repeat application at 7-10 day intervals. **NOTE:** Sulfur may injure certain varieties of beans. For control of Powdery and Downy Mildew and Botrytis Blight on **Onions.** Begin application before diseases are expected to appear. Repeat as necessary.

FRUIT TREES: (As Fungicide & Insecticide)

APPLE, PEAR: For Powdery Mildew and Scab. Reduce dosage to 1 tablespoon per gallon of water if hot weather threatens. For disease control on apples apply from prepink through cover periods; for pears apply at prebloom, bloom, petal fall, and cover periods.

PEACH: For Brown Rot, Scab. Apply in pink, bloom, shuck, cover and pre-harvest sprays.

CHERRY: For Brown Rot, Leaf Spot. Apply in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

PLUM, PRUNE: For Brown Rot, Leaf Spot. Apply in pink, bloom, shuck, cover and pre-harvest sprays.

CITRUS: For Brown Rot. Apply according to state recommendations. For Rust Mite, Thrips, Red Spider Mites, Two-Spotted Spider Mites. Begin applications at first signs of damage and repeat as necessary. Rust mites should be controlled at all times of the year in order to prevent russeting of fruit and greasy spots on leaves.

COMMON CHIGGER CONTROL: For control of chiggers in the yard, dust over the entire lawn, flower garden, shrubs and any place the chiggers may be hiding. This should be done a day or two before protection is desired and should be repeated at weekly intervals. The contents of this package will cover from 1500 to 2000 square feet. Dust on clothing especially around the legs when visiting chigger-infested areas.

3. HERE ARE THE RESULTS YOU MAY EXPECT

Efficient and economical control of listed diseases and insects may be had with use of this product.

CAUTION: Keep out of reach of children. Do not get in eyes. Do not use within 2 weeks of an oil spray treatment, on Citrus within 3 to 6 weeks. Use only according to state recommendations. Sulfur may burn foliage when temperature is high (90-95°F.). Do not make applications at such times. Do not use on cucurbits (Cucumbers, Cantaloupe, Melons, Squash) or Tung.

Do not reuse empty container. Destroy it by burying with waste or burning. Stay away from smoke or fumes.

Buyer assumes all risk of use, storage or handling of this material not in strict accordance with directions given herewith.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS