

1003
010-1
7401-180

(Front Panel)

ferti-lome(R)

ACCEPTED
MAR 24 1987
Under the Federal Fungicide, and Insecticide Act, as amended, for the use registered under EPA Reg. No. 7401-180

Dusting

Sulfur

Use as a Dust or Spray!

- Controls Black Spot, Powdery Mildew, Leaf Spot, Rust & Brown Canker on Roses.
- Controls Listed Diseases & Insects on Shrubs, Flowers, Vegetables & Fruits.
- Controls Thrips, Rust Mites, Red Spider Mites & Two Spotted Mites on Citrus.
- Controls Chiggers

Active Ingredient:
 Sulfur..... 90%
 Inert Ingredients:..... 10%

KEEP OUT OF REACH OF CHILDREN (12 pt)

CAUTION (18 pt)

See Back Panel For Additional Precautionary Statements

(Back Panel)

DIRECTIONS FOR USE: It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

1. WHAT IS THIS PACKAGE FOR?

This product provides effective and economical control of listed plant diseases and insects. May be used as a dust or spray.

2. HOW DO YOU USE IT?

When used as a dust, punch out perforated tab on top of shaker can and rotate lid to dust setting. Apply to all plant surfaces to thoroughly cover. When used as a spray, mix 4 tablespoons per each 1 gallon of water used. Spray to cover thoroughly all plant surfaces. NOTE: DO NOT RE-ENTER TREATED AREAS FOR AT LEAST 24 HOURS AFTER APPLICATION IS MADE.

ROSES, SHRUBS, FLOWERS: (as a Fungicide & Insecticide)

For Black Spot, Brown Canker, Leaf Spot, Powdery Mildew, Thrips and Mites on Roses. Begin spraying when new growth starts in the spring. Repeat at 7 day intervals and re-apply after rainfall. For disease and insect control on shrubs, flowers and ornamentals begin application when diseases appear and repeat at 5-10 day intervals and after rains.

For control of Powdery Mildew, Thrips and Mites on Azalea, Alder, Buttonbush, Catalpa, Cosmos, Crape Myrtle, Dahlia, Daisy, Delphinium, Aster, Calendula, Dogwood, Eucnymus, Gladiolus, Golden Fleece, Hibiscus, Holly, Hydrangeas, Lady's Mantel, Ligustrum, Lilac, Petunia, Poplars, Silver Vine, Spirea, Sunflower, Sweet Pea, Verbena, Violets, Willows, Carnations, Cedar, Cherry Laurel, Chrysanthemums, Hollyhock, Juniper, Simlax and Spruce, For control of Leaf Spot, Thrips, and Mites on Calendula, Cedar, Cherry Laurel, Chrysanthemum, Clematis, Columbine, Foxglove, Hollyhock, English Ivy, Juniper, Laurel, Sage, Simlax, Snapdragons, and Spruce. For control of Rust, Thrips, and Mites on Asters, Carnations, Chrysanthemums, Hollyhock and Snapdragon.

VEGETABLES: (as Fungicide)

For control of Powdery Mildew on Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower. For control of Rust on Beans and Peas. Begin applications at first sign of disease. Repeat applications at 7-10 day intervals. NOTE: Sulfur may injure certain varieties of beans. For control of Powdery and Downy Mildew and Botrytis Blight on Onions. Begin application before diseases are expected to appear. Repeat as necessary. Do not use on Cucurbits (Cucumbers, Cantaloupes, Melons, Squash) or Tung.

FRUITS: (as Fungicide and Insecticide)

APPLE, PEAR: For Powdery Mildew and Scab. Reduce dosage to 1 tablespoon per gallon of water if hot weather threatens. For disease control on apples apply from pre-pink through cover periods. For Pears apply at prebloom, bloom, petal fall and cover periods.

PEACH: For Brown Rot, Scab. Apply in pink, bloom, shuck, cover and preharvest sprays.

CHERRY: For Brown Rot, Leaf Spot. Apply in pink, bloom, petal fall, shuck, cover and post-harvest sprays.

PLUM, PRUNE: For Brown Rot, Leaf Spot. Apply in pink, bloom, shuck, cover and pre-harvest sprays.

CITRUS: For Brown Rot. Apply according to state recommendations. For Rust, Mite, Thrips, Red Spider Mites, Two-spotted Spider Mites. Begin applications at first signs of damage and repeat as necessary. Rust Mites should be controlled at all times of the year in order to prevent russetting of fruit and greasy spots on leaves.

BEST AVAILABLE COPY

2003

(Back Panel Cont'd)

COMMON CHIGGER CONTROL: For control of chiggers in the yard, dust over the entire lawn, flower garden, shrubs and any place the chiggers may be hiding. This should be done a day or two before protection is desired and should be repeated at weekly intervals. The contents of this package will cover from 1,500 to 2,000 square feet.

IMPORTANT: Do not use within 2 weeks of an oil spray treatment, on citrus within 3 to 6 weeks. Use only according to state recommendations. Sulfur may burn foliage when temperature is high (90-95 F).

3. HERE ARE THE RESULTS YOU MAY EXPECT!

When used as directed, this product will provide effective and economical control of listed diseases and insects and thus promote healthy, lush growth.

DISPOSAL (12 pt)

Do not reuse empty container. Wrap container and put in trash collection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION: May be harmful if swallowed, inhaled or in eyes. Avoid contact with skin, eyes or clothing. Wash thoroughly after handling and before eating or smoking. Keep children and pets off treated areas for at least 24 hours and until this material has washed into the soil.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES, flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN, wash affected areas with plenty of soap and water.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water.

Buyer assumes all risks of use, storage, or handling of this material not in strict accordance with directions given herewith. Voluntary Purchasing Groups, Inc. warrants that this material conforms to the chemical description on the label and suited for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

Voluntary Purchasing Groups, Inc.
Eonham, Texas 75418

EPA Est. No. 7401-TX-1

EPA Reg. No. 7401-188

Net Contents 2 lbs.

BEST AVAILABLE COPY