

U.S. ENVIRONMENTAL PROTECTION AGENCY

Office of Pesticide Programs Registration Division (7505P) 1200 Pennsylvania Ave., N.W. Washington, D.C. 20460

7401-188

EPA Reg. Number:

Date of Issuance:

5/31/2018

NOTICE OF PESTICIDE:

X Registration
X Reregistration
(under FIFRA, as amended)

Term of Issuance: Unconditional

Name of Pesticide Product:

Ferti-Lome Dusting Sulfur

Name and Address of Registrant (include ZIP Code):

Ann M. Tillman Agent for Voluntary Purchasing Groups, Inc. 230 FM 87

Bonham, TX 75418

Note: Changes in labeling differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby registered/reregistered under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended. Registration is in no way to be construed as an endorsement or recommendation of this product by the Agency. In order to protect health and the environment, the Administrator, on her/his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

EPA received a label amendment request submitted by email on 03/21/2018. EPA grants this request under the authority of section 3(c)(5) of FIFRA, as amended. With this accepted labeling, all requirements set forth in the Reregistration Eligibility Decision for Sulfur have been satisfied. Therefore, EPA reregisters the product listed above. This action is taken under the authority of section 4(g)(2)(c) of FIFRA, as amended. Reregistration under this section does not eliminate the need for continual reassessment of pesticides. EPA may require submission of data at any time to maintain the registration of your product.

Submit one (1) copy of final printed labeling. Amended labeling will supersede all previously accepted labels. A copy of your label stamped "Accepted" is enclosed for your records. Products shipped after 12 months from the date of this Notice or the next printing of your label, whichever occurs first, must bear the new revised label.

Signature of Approving Official:	Date:
Erik Kraft, Product Manager 24 Fungicide & Herbicide Branch, Registration Division (7505P)	5/31/2018

EPA Form 8570-6

Page 2 of 2 EPA Reg. No. 7401-188 Decision No. 318063

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6. Your release for shipment of the product constitutes acceptance of these conditions. A stamped copy of the label is enclosed for your records.

Please also note that the record for this product currently contains the following CSFs:

• Basic CSF dated 04/23/2018

If you have any questions, please contact Lisa Pahel by phone at (703) 347-0459, or via email at pahel.lisa@epa.gov.

{Note to reviewer: brackets {} indicate notes to reviewer or location of text on label} [] indicates alternate text {FRONT LABEL PANEL}

Ferti-lome® Dusting Sulfur [Alternate Brand Name Snake Eyes Brand Dusting Sulfur]





SULFUR GROUP M02 FUNGICIDE

[Optional proposed text appears in brackets-the final label may include some or all of the optional text]

- [For Use in Residential Lawns and Gardens]
- [Controls Black Spot, Powdery Mildew, Leaf Spot, Rust & Brown Canker on Roses.]
- [Controls Listed Diseases & Insects on Shrubs, Flowers, Vegetables & Fruits]
- [Controls Thrips, Rust Mites, Red Spider Mites & Two Spotted Mites on Citrus.]

ACTIVE INGREDIENT:

Sulfur	90.0%
INERT INGREDIENTS:	10.0%
TOTAL:	100.0%

ACCEPTED

05/31/2018

Under the Federal Insecticide, Fungicide and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No.

7401-188

KEEP OUT OF REACH OF CHILDREN

CAUTION

See Back/Side Panel for Additional Precautionary Statements

Net Weight

{BACK/SIDE LABEL PANEL}

FIRST AID	
If inhaled	 Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	 Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	 Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	 Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful absorbed through skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wear long pants, long sleeved shirt, shoes, socks, water resistant gloves, and goggles or face shield. Wash thoroughly after handling and before eating or smoking.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Keep children and pets off treated areas for at least 24 hours and until this material has washed into the soil.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes and being grown in home lawns and gardens.

{BACK/SIDE LABEL PANEL}

1. WHAT IS THIS PACKAGE FOR?

This product provides effective and economical control of listed plant diseases and insects. May be used as a dust or spray.

2. HOW DO YOU USE IT?

When used as a dust, punch out perforated tab on top of shaker can and rotate lid to dust setting. Apply to all plant surfaces to thoroughly cover. When used as a spray, mix 4 tablespoons (1.33 oz.) per each 1 gallon of water used (0.0832 lb/gallon). Spray to cover thoroughly all plant surfaces. NOTE: DO NOT RE-ENTER TREATED AREAS FOR AT LEAST 24 HOURS AFTER APPLICATION IS MADE.

ROSES, SHRUBS, FLOWERS: (as a Fungicide & Insecticide)

For Black Spot, Brown Canker, Leaf Spot, Powdery Mildew, Thrips and Mites on Roses. Begin spraying when new growth starts in the spring. One gallon covers 35 sq. ft. Repeat applications at 7 to 10 day intervals and re-apply after rainfall. For disease and insect control on shrubs, flowers and ornamentals begin application when diseases appear and repeat at 5-10 day intervals and after rains. For control of Powdery Mildew, Thrips and Mites on Azalea, Alder, Buttonbush, Catalpa, Cosmos, Crape Myrtle, Dahlia, Daisy, Delphinium, Aster, Calendula, Dogwood, Euonymus, Gladiolus, Golden Fleece, Hibiscus, Holly, Hydrangeas, Lady's Mantel, Ligustrum, Lilac, Petunia, Poplars, Silver Vine, Spires, Sunflower. Sweet Pea, Verbena, Violets, Willows, Carnations, Cedar, Cherry Laurel, Chrysanthemums, Hollyhock, Juniper, Simlax and Spruce. For control of Leaf Spot, Thrips, and Mites on Calendula, Cedar, Cherry Laurel, Chrysanthemum, Clematis, Columbine., Foxglove, Hollyhock, English Ivy, Juniper, Laurel, Sage, Simlax, Snapdragons, and Spruce. For control of Rust, Thrips, and Mites on Asters, Carnations, Chrysanthemums, Hollyhock and Snapdragon.

VEGETABLES: (as Fungicide)

For control of Powdery Mildew on Beans, Peas, Broccoli, Brussels Sprouts, Cabbage, Cauliflower. For control of Rust on Beans and Peas. One gallon covers 100 to 1,000 sq. ft. Apply one gallon per 100 sq. ft. during periods of frequent and heavy rainfall. Begin applications at first sign of disease. Repeat applications at 7-10 day intervals. NOTE: Sulfur may injure certain varieties of beans. For control of Powdery and Downy Mildew and Botrvtis Blight on Onions. Begin application before diseases are expected to appear. Repeat up to the day of harvest.

Restrictions: Do not use on Cucurbits (Cucumbers, Cantaloupes, Melons, and Squash) or Tung.

FRUITS:(as Fungicide and Insecticide)

APPLE, PEAR: For Powdery Mildew and Scab. Reduce dosage to 1 tablespoon (0.33 oz) per gallon of water if hot weather threatens. One gallon covers 60-360 sq. ft. Apply one gallon per 60 sq. ft. during periods of frequent and heavy rainfall. For disease control on apples apply from pre-pink through cover periods. For Pears apply at pre-bloom, bloom, petal fall and cover periods. Repeat applications every 7 to 14 days as needed up to day of harvest.

{BACK/SIDE LABEL PANEL}

PEACH: For Brown Rot, Scab. Apply in pink, bloom, shuck, cover and pre-harvest sprays. One gallon covers 75-90 sq. ft. Apply one gallon per 75 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 10 to 14 days as needed up to day of harvest.

CHERRY: For Brown Rot, Leaf Spot. Apply in pink, bloom, petal fall, shuck, cover and post-harvest sprays. One gallon covers 70 to 180 sq. ft. Apply one gallon per 70 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 7 to 14 days as needed up to day of harvest.

PLUM, PRUNE: For Brown Rot, Leaf Spot. Apply at pink, bloom, shuck, cover and pre-harvest sprays up to day of harvest. One gallon covers 75-90 sq. ft. Apply one gallon per 75 sq. ft. during periods of frequent and heavy rainfall. Repeat applications every 7 to 14 days as needed up to day of harvest.

CITRUS: For Brown Rot. Apply according to state directions up to day of harvest. For Rust, Mite Thrips, Red Spider Mites, Two-spotted Spider Mites. Begin applications at first signs of damage and repeat as needed. Rust Mites should be controlled at all times of the year in order to prevent russeting of fruit and greasy spots on leaves.

Restriction: Do not use within 2 weeks of an oil spray treatment, on citrus within 3 to 6 weeks. Use only according to state directions. Sulfur may burn foliage when temperature is high (90-95 F).

3. HERE ARE THE RESULTS YOU MAY EXPECT!

When used as directed, this product will provide effective and economical control of listed diseases and insects and thus promote healthy, lush growth.

STORAGE AND DISPOSAL

PESTICIDE STORAGE: Keep out of reach of children and pets. Store in original containers only. Store in cool, dry place and avoid excess heat.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Non-refillable container. Do not reuse this container. **If Empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith. To the extent consistent with applicable law, Voluntary Purchasing Groups, Inc. warrants this product to conform to the chemical description on the label and for the purposes stated on the label. The limit of any liability incurred shall be the purchase price paid by the user or buyer.

[optional bar code]

Manufactured By:
VOLUNTARY PURCHASING GROUPS, INC.
230 FM 87
BONHAM, TEXAS 75418
EPA Reg. No. 7401-188
EPA Est. No. 7401-TX-1
www.fertilome.com