

Center Panel  
VPG  
HERBAN 62  
DECIMATE<sub>R</sub> + HERBAN\*\* SURFACTANT

ACTIVE INGREDIENTS:	
norea (3-(hexahydro-4, 7-methanoindan-5-yl)-1,1-dimethylurea)*....	13.1%
related reaction products .....	0.7%
Monosodium Acid Methanearsonate .....	29.0%
INERT INGREDIENTS: .....	57.2%
Total .....	100.0%

Total arsenic content, as elemental, all in water soluble form ---- 13.3%  
Product contains 3.3 lbs. MSMA per gallon, and 1.6 lbs. norea and related  
reaction products per gallon.

\*Herban is the registered trade mark of Hercules Powder Co.

CAUTION - - - 18 point type

Keep Out Of Reach Of Children - - - 14 point type

See Side Panel For Additional Cautions

Sold By


Voluntary Purchasing Groups, Inc.

Bonham, Texas

Net Contents One Gallon

Right Hand Panel

SHAKE WELL BEFORE MIXING OR SAMPLING



CAUTION: Harmful if swallowed. Avoid contact with skin. Avoid breathing  
spray mist. Wash hands after using. Avoid storage near feed or food pro-  
ducts. Keep away from children and domestic animals. Do not get in eyes.

Do not contaminate waters used by fish and wildlife, domestic animals, or  
for irrigation or domestic water supplies.  
Keep children and animals off treated areas until washed into soil.

Do not graze treated areas or feed livestock treated forage.

Following Crops - Herban is gradually destroyed in the soil. However, crops  
other than cotton, sorghum, seed spinach and sugar cane should not be planted  
sooner than 5 months after treatment of soil with herban to avoid the possi-  
bility of crop injury.

Destroy container - Do not re-use for any other purpose.

Prevent Injury to other crops or Plants - Do not use herban-MSMA on or near  
lawns, flowers, or other desirable plants. Prevent the drift of spray to  
other areas.

NON-WARRANTY: Seller warrants that the product conforms to its chemical  
description and is reasonably fit for the purpose stated on the label  
when used in accordance with directions under normal conditions of use, but  
neither this warranty nor any other warranty of merchantability or fitness for  
a particular purpose, express or implied, extends to the use of this product  
contrary to label instructions, or under abnormal conditions, or under con-  
ditions not reasonably foreseeable to seller, and buyer assumes the risk of  
any such use.

Left Hand Panel

SHAKE WELL BEFORE MIXING OR SAMPLING

DIRECTIONS FOR USE:

Equipment - A sprayer designed for low volume application is recommended. Do not use strainers finer than 50 mesh. The spray mixture must be agitated to maintain a uniform spray concentration. By-pass agitation is satisfactory.

Application of the Spray - 30 to 40 gallons of spray broadcast rate per acre are required to ensure thorough uniform coverage of weeds. Use a proportional amount for band treatment depending on band width treated. Keep off cotton foliage, direct spray on weeds.

Mixing the Spray - Shake well before mixing. Add about  $\frac{1}{2}$  of the total volume of water to the tank. Start the pump to provide agitation, and pour in the desired amount of herbicide concentrate. Rinse herbicide containers into spray tank. Add water to make the total desired volume.

Accurate Dosage - Calibrate equipment and double check calculations and measurements for preparation of spray liquid. Shut off spray when turning, stopping or slowing tractor.

Pre-Bloom, Post-Emergence Weed Control in Cotton - For broad spectrum post-emergence weed control, the Herban-MSMA-Surfactant mix may be used to control Johnson grass, Goose grass, Nutgrass, (Coco), Colorado grass, Barnyard grass (Water grass), Crabgrass, Morning glory, Cocklebur, Pigweed, Purslane, Coffee Weed, Florida pusley, Smartweed, Poverty weed and Carpetweed.

Rates - Use  $\frac{1}{2}$  gallon broadcast rate in 30 to 40 gallons of water per acre. If band application is used, apply a proportional amount of the mixture. For example, on a 14" band (40" row spacing) use 10-14 gallons of spray mixture per acre of crop. Applications should be made to weeds when cotton is 3 inches high and until first bloom. Do not apply this mixture after the first bloom opens. Thoroughly wet stems and foliage of weeds. Nozzles should be arranged so as to avoid spraying the cotton foliage but also to give maximum coverage of weeds and grasses. Make a second application at the same rate 7 days after the first. Do not add additional wetting agent to the spray mixture.

USDA Reg. No. 4272-83

DO NOT USE OR STORE IN OR AROUND THE HOME

---- 18 point type

DO NOT ALLOW DOMESTIC ANIMALS TO GRAZE TREATED AREA

