

	Center Label	
1-1-1	1-YIELD	4.00
	DECI-MEK	1.00
	PLURIPAC	
AC 14-1-1		
Arsenic acid		1.00
Monosodium Arsenite		15.00
Diethylarsinic acid		44.00
	total	60.00
total amount of water soluble form		20.00

POISON POISON

DANGER - - - 10 point type

keep out of reach of children

- - - - 14 point type

see side panel for Additional Cautions & Antidote

ACCEPTED

DEC 13 1967

UNDER THE FEDERAL INSECTICIDE,
FUNGICIDE AND ROENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
UNDER NO. 272-28 SUBJECT
TO ATTACHED COMMENTS.

7401-20

1-1-1

Houston, Texas

net contents 1 gallon

side panel

WARNING

Avoid contact with skin, eyes, or clothes. Do not breathe mist; wear respirator. Wash thoroughly after using and before eating or smoking. Avoid contamination of food and water to be consumed by humans and animals. Keep livestock, poultry, pets and children off of treated areas. Do not graze or use treated plants for food or forage. Since 1-Yield 101 is injurious to all plant foliage, avoid spray drift to areas and buildings occupied by humans, animals or poultry.

Do not contaminate streams, lakes, ponds, irrigation water, wetlands or similar areas.

Do not apply over root zone of desirable plants.

WASH EQUIPMENT: Wash spray equipment and containers and dispose of liquid wastes in a pit in non-crop land located away from water supplies.

Right Hand Panel
(Cont'd)



POISON



ACTION:

INGESTION: Call a physician immediately. Give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Then give 2 tablespoonfuls of Epsom Salts or Milk of Magnesia in water. Have victim lie down and keep quiet.

SKIN CONTACT: In case of spillage on skin, wash promptly with soap and water.

Product No. 4272-

Left Hand Panel

DIRECTIONS FOR USE

For weed control on industrial sites, driveways, implement storage yards, drainage ditchbanks, and other non-crop areas, mix 2 quarts in sufficient water to cover 1 acre (10-15 gallons). Apply with low pressure ground spray equipment (about 25 lbs. pressure). Adequate coverage is essential for good weed control. Best control is obtained when temperature averages 70 degrees or above.

This application will control goose grass, Cocklebur, Green and Yellow Fox-tail, Pigweed, Milkweed, Crabgrass, Lambsquarter, Larkspur, Ragweed, and give top kill of Nutgrass and Johnson grass. Repeat application may be necessary to control weeds not properly covered with spray in from 7-10 days. Apply when weeds are young.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION:

i-Yield 200 has acidic properties and reacts with fabric, galvanized metals, black iron, and certain other metals. Any metallic equipment or nozzles used in application of i-Yield 200 should be made of stainless steel, as there is little or no reaction with this metal at field strength dilution. Do not use galvanized metal containers or spraying equipment as highly toxic arsine gas may be formed. To reduce possible corrosion to metals and deterioration of other parts, all equipment which has been in contact with or exposed to i-Yield 200 or its solution, should be thoroughly washed with water immediately after use. i-Yield 200 should not be stored in tightly sealed, unlined metal containers because of the possibility of excessive pressure developing from hydrogen liberation.

DO NOT USE OR STORE IN Airtight Containers - - - In point type

DO NOT ALLOW DRAINAGE TO A PLANT TO BE OPEN TO AIR - - - In point type

HI-YIELD DECIMEX PLUS SURFACTANT 4272-R

This product contains:

2.39 lbs. MSMA per gallon

5.06 lbs. Arsenic Acid per gallon