

(Front Panel)

HI-YIELD

SEVIN* EMULSIFIABLE CONCENTRATE

(Contains 2 lbs. Carbaryl per gallon)

ACTIVE INGREDIENT:

Carbaryl: (1-naphthyl N-Methylcarbamate)*	25.00%
INERT INGREDIENTS:	75.00%

Total . . 100.00%

*Sevin is the registered trademark of Union Carbide Corporation for Carbaryl Insecticide

CAUTION! (18 Point)

KEEP OUT OF REACH OF CHILDREN (12 Point)

SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

Harmful if Inhaled or Swallowed. Avoid Breathing of Spray Mist. Skin Contact may be harmful.

Avoid contact. Take shower or bathe after work. Wear regular long sleeved work clothing. Change to clean clothing daily.

Wash hands and face before eating.

Wash thoroughly after handling.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES AND PONDS.

DO NOT USE, POUR, SPILL OR STORE NEAR HEAT OR OPEN FLAME.

DO NOT REUSE CONTAINER FOR ANY PURPOSE. DESTROY EMPTY CONTAINER.

VOLUNTARY PURCHASING GROUPS, INC.

BONHAM, TEXAS

NET CONTENTS GALLONS

ACCEPTED

AUG 23 1972

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POLICE REGISTER-
ED UNDER NO. SUBJECT
TO ATTACHED COMMENTS.

7401-~~210~~ 210

4272-99

(Side Panel)

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

NOTE

1. Compatible with commonly used insecticides such as lead arsenate, Methoxychlor and other chlorinated hydrocarbons; Guthion, Malathion, Parathion and other organic phosphates and botanicals; also with such fungicides as glyodin, fixed coppers, dithiocarbamates, sulfur, captan, Cyprex and Phaltan.
2. Unstable under highly alkaline conditions.
3. Some phytotoxicity may occur on tender foliage in presence of rain or high humidity of several days duration following spraying.
4. Does not control spider mites.
5. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.
6. Allow 7 days before grazing or harvesting crop residue for feed or bedding for dairy or meat animals. Tolerance 100 ppm or forage.

NON-WARRANTY

Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

GENERAL DIRECTIONS

RECOMMENDED SPRAYING CONCENTRATION OF 10% SEVIN* EMULSIFIABLE CONCENTRATE in hydraulic sprayers, low volume ground applicators or airplane sprayers.

LOW VOLUME GROUND EQUIPMENT: Apply at least 5 gallons per acre of prepared spray. Insure uniform coverage.

AIRPLANE APPLICATION: Apply 3-10 gallons per acre of prepared spray.

MIXING DIRECTIONS

Mix with water by agitation; agitate the mixture while spraying. If allowed to stand, re-agitate before use.

DIRECTIONS FOR USE

Recommended dosages refer to quarts of SEVIN* EMULSIFIABLE CONCENTRATE per acre.

COTTON INSECT CONTROL

Use 1-2 quarts for early-season thrips, flea beetles, cotton leafhoppers, striped blister beetle and cotton leafworm. Use 2-3 quarts, depending on the size of the cotton and insect population level, for mid- and late-season boll weevil, bollworms, fall armyworms, cotton leaf perforator, leaf rollers, cotton fleahoppers and tarnished plant bug. Also for light-to-moderate infestations of Lygus bug in Western irrigated cotton.

Applications should be made every 5-7 days as long as control is necessary.

Use 3-5 quarts for pink bollworm on a 5-7 day schedule depending on the size of cotton and degree of infestation. Use 1-3 quarts for grasshoppers. The highest dosage should be used when grasshoppers are mature or material is applied to partly defoliated plants. Aphid populations will be suppressed by repeated applications of this insecticide. This insecticide does not control spider mites. When infestations are encountered use a recommended miticide.

SOYBEAN INSECT CONTROL

Use 1-2 quarts for blister beetle and Mexican bean beetle.

Use 2 quarts for alfalfa caterpillar, bean leaf beetle, cucumber beetle, green cloverworm, Japanese beetle, leafhoppers, three-cornered alfalfa hopper, thrips and velvet bean caterpillar. Use 2-3 quarts for armyworms, corn earworm (bollworm), stink bugs and webworms.

To avoid possible injury on tender foliage, do not apply when foliage is wet or when rain or excessive humidity is expected during the next 2 days.

TOBACCO INSECT CONTROL

Use 2-4 quarts for budworm, flea beetle, Japanese beetle, June beetle and hornworm. Use lower rate on young plants up to knee high; higher rate for mature plants. Begin treatment when worms are small. Avoid excessive application in plant bed as injury may result. Treated fields may be entered immediately after foliage has dried. If late applications are necessary, allow 3 days before priming or cutting.

VEGETABLE INSECT CONTROL

Recommended dosages refer to quarts of SEVIN* EMULSIFIABLE CONCEN per acre. When a dose range is indicated use lower rate on young plants; higher rate on mature plants. Use sufficient spray volume to obtain full coverage. Keep spray mixtures well agitated. Apply when insects or their damage appear. Repeat at 7-10 day intervals or as necessary unless shorter interval is specified below. On vegetables intervals in days between application and harvest of crop that must be observed are indicated in () parenthesis after each crop heading.

GARDEN PEAS (0) Use 2 quarts for leafhoppers, Colorado potato beetle. For Colorado potato beetle control apply to weeds in fields of garden peas to help prevent insect contamination in processed peas. Use 2-3 quarts for armyworms and grasshoppers. Repeat as necessary.

POTATO (0), TOMATO (0), EGG PLANT (0) and PEPPER (0): Use 1-2 quarts for Colorado potato beetle, flea beetles and leafhoppers. Use 2-4 quarts for European corn borer, fall armyworm, lace bugs, tomato fruitworm, tomato hornworm, tarnished plant bugs and stink bugs. Use 4 quarts for cutworm control on potato, tomato and eggplant.

OKRA (0): Use 2-4 quarts for corn earworm and stink bugs. Repeat at 5-7 day intervals or as necessary.

CABBAGE (3), BROCCOLI (3), BRUSSELS SPROUTS (3), CAULIFLOWER (3)

bug. Use 2-4 quarts for armyworm, imported cabbage worm and corn earworms. When cabbage looper is the principal problem use an alternative pesticide that is specifically recommended. For control of small cabbage loopers in home gardens or in commercial plantings after edible portions begin to form, use Sevin in a 5-7 day schedule.